

(19)



Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11) Publication number:

**0 534 363 A2**

(12)

## EUROPEAN PATENT APPLICATION

(21) Application number: **92116180.8**

(51) Int. Cl. 5: **C07K 5/06, A61K 37/64**

(22) Date of filing: **21.09.92**

(30) Priority: **27.09.91 US 767286**  
**20.08.92 US 929481**

(43) Date of publication of application:  
**31.03.93 Bulletin 93/13**

(84) Designated Contracting States:  
**AT BE CH DE DK ES FR GB GR IE IT LI LU MC**  
**NL PT SE**

(71) Applicant: **MERRELL DOW**  
**PHARMACEUTICALS INC.**  
**2110 East Galbraith Road**  
**Cincinnati Ohio 45215(US)**

(72) Inventor: **Warshawsky, Alan M.**  
**10713 Shadowcrest Ct.**  
**Cincinnati, Ohio 45242(US)**  
Inventor: **Flynn, Gary A.**  
**7121 Euclid Road**  
**Cincinnati, Ohio 45243(US)**

(74) Representative: **Vossius & Partner**  
**Siebertstrasse 4 P.O. Box 86 07 67**  
**W-8000 München 80 (DE)**

(54) **Novel 2-substituted Indane-2-mercaptoacetylamine derivatives useful as inhibitors of**  
**enkephalinase and ace.**

(57) The present invention relates to novel 2-substituted indane-2-mercaptoacetylamine derivatives which are useful as inhibitors of enkephalinase and ACE.

**0 534 363 A2**

This application is a continuation-in-part of application Serial No. 07/767,286, filed September 27, 1991.

Enkephalinase or, more specifically, endopeptidase-24.11, is a mammalian ectoenzyme which is involved in the metabolic degradation of certain circulating regulatory peptides. This enzyme, which is a  $Zn^{+2}$ -metallopeptidase, exerts its effect by cleaving the extracellular peptides at the amino group of hydrophobic residues and thus inactivates the peptides as regulatory messengers.

Enkephalinase is involved in the metabolic degradation of a variety of circulating regulatory peptides including endorphins, such as  $\beta$ -endorphin and the enkephalins, atrial natriuretic peptide (ANP), and other circulating regulatory peptides.


Endorphins are naturally-occurring polypeptides which bind to opiate receptors in various areas of the brain and thereby provide an analgesic effect by raising the pain threshold. Endorphins occur in various forms including  $\alpha$ -endorphin,  $\beta$ -endorphin,  $\gamma$ -endorphin as well as the enkephalins. The enkephalins, i.e., Met-enkephalin and Leu-enkephalin, are pentapeptides which occur in nerve endings of brain tissue, spinal cord and the gastrointestinal tract. Like the other endorphins, the enkephalins provide an analgesic effect by binding to the opiate receptors in the brain. By inhibiting enkephalinase, the metabolic degradation of the naturally-occurring endorphins and enkephalins are inhibited, thereby providing a potent endorphin- or enkephalin-mediated analgesic effect. Inhibition of enkephalinase would therefore be useful in a patient suffering from acute or chronic pain. Inhibition of enkephalinase would also be useful in providing an antidepressant effect and in providing a reduction in severity of withdrawal symptoms associated with termination of opiate or morphine administration.

ANP refers to a family of naturally-occurring peptides which are involved in the homeostatic regulation of blood pressure, as well as sodium and water levels. ANPs have been found to vary in length from about 21 to about 126 amino acids with a common structural feature being one or more disulfide-looped sequences of 17 amino acids with various amino- and carboxy-terminal sequences attached to the cystine moiety. ANPs have been found to bind to specific binding sites in various tissues including kidney, adrenal, aorta, and vascular smooth muscle with affinities ranging from about 50 pico-molar (pM) to about 500 nano-molar (nM) [Needleman, *Hypertension* 7, 469 (1985)]. In addition, it is believed that ANPs bind to specific receptors in the brain and possibly serve as a neuromodulator as well as a conventional peripheral hormones.

The biological properties of ANP involve potent diuretic/natriuretic and vasodilatory/hypotensive effects as well as an inhibitory effect on renin and aldosterone secretion [deBold, *Science* 230, 767 (1985)]. By inhibiting enkephalinase, the metabolic degradation of the naturally-occurring ANP is inhibited, thereby providing a potent ANP-mediated diuretic, natriuretic, hypotensive, hypoaldosteronemic effects. Inhibition of enkephalinase would therefore be useful in a patient suffering from disease states characterized by abnormalities in fluid, electrolyte, blood pressure, intraocular pressure, renin, or aldosterone homeostasis, such as, but not limited to, hypertension, renal diseases, hyperaldosteronemia, cardiac hypertrophy, glaucoma and congestive heart failure.

In addition, the compounds of the present invention are inhibitors of Angiotensin-Converting Enzyme (ACE). ACE is a peptidyl dipeptidase which catalyzes the conversion of angiotensin I to angiotensin II. Angiotensin II is a vasoconstrictor which also stimulates aldosterone secretion by the adrenal cortex. Inhibition of ACE would therefore be useful in a patient suffering from disease states such as hypertension and congestive heart failure [See William W. Douglas, "Polypeptides - Angiotensin, Plasma Kinins, and Others", Chapter 27, in GOODMAN AND GILLMAN'S THE PHARMACOLOGICAL BASIS OF THERAPEUTICS, 7th edition, 1985, pp. 652-3, MacMillan Publishing Co., New York, New York]. In addition, it has been disclosed that ACE inhibitors are useful in treating Cognitive disorders [German Application No. 3901-291-A, published August 3, 1989].

The present invention provides novel compounds of the Formula (I)



wherein

B<sub>1</sub> and B<sub>2</sub> are each independently hydrogen; hydroxy; -OR<sub>2</sub> wherein R<sub>2</sub> is a C<sub>1</sub>-C<sub>4</sub> alkyl or an Ar-Y group wherein Ar is aryl and Y is a hydrogen or C<sub>1</sub>-C<sub>4</sub> alkyl; or, where B<sub>1</sub> and B<sub>2</sub> are attached to adjacent carbon atoms, B<sub>1</sub> and B<sub>2</sub> can be taken together with said adjacent carbons to form a benzene ring or methylenedioxy;


A is a bond, methylene, oxygen, sulfur,  $\text{NR}_4$  or  $\text{NCOR}_5$  wherein  $\text{R}_4$  is hydrogen, a  $\text{C}_1$ - $\text{C}_4$  alkyl or an Ar-Y-group and  $\text{R}_5$  is  $-\text{CF}_3$  or a  $\text{C}_1$ - $\text{C}_{10}$  alkyl or an Ar-Y-group;

R<sub>1</sub> is hydrogen, acetyl, -CH<sub>2</sub>OC(O)C(CH<sub>3</sub>)<sub>3</sub> or benzoyl;

$R_3$  is hydrogen or  $-\text{CH}_2\text{OC}(\text{O})\text{C}(\text{CH}_3)_3$ ;

n is an integer 0 to 3; and

**Q** is a group of the formula



wherein Z is O, NH or S; and m is an integer 1 to 5.

The present invention further provides a compound of Formula I for use in inhibiting enkephalinase in a patient in need thereof comprising administering to said patient an effective enkephalinase inhibitory amount of a compound of Formula (I). The present invention also provides a compound of Formula I for use in inhibiting ACE in a patient in need thereof comprising administering to said patient an effective ACE inhibitory amount of a compound of Formula (I).

In addition, the present invention provides a composition comprising an assayable amount of a compound of Formula (I) in admixture or otherwise in association with an inert carrier. The present invention also provides a pharmaceutical composition comprising an effective inhibitory amount of a compound of Formula (I) in admixture or otherwise in association with one or more pharmaceutically acceptable carriers or excipients.

As used herein, the term "C<sub>1</sub>-C<sub>4</sub> alkyl" refers to a saturated straight or branched chain hydrocarbon radical of one to four carbon atoms and includes methyl, ethyl, propyl, isopropyl, n-butyl, isobutyl, tertiary butyl and the like. The terms "C<sub>1</sub>-C<sub>8</sub> alkyl" and "C<sub>1</sub>-C<sub>10</sub> alkyl" refer to saturated straight or branched chain hydrocarbon radicals of one to eight and one to ten carbon atoms, respectively, including methyl, ethyl, propyl, isopropyl, n-butyl, isobutyl, tertiary butyl, pentyl, isopentyl, hexyl, 2,3-dimethyl-2-butyl, heptyl, 2,2-dimethyl-3-pentyl, 2-methyl-2-hexyl, octyl, 4-methyl-3-heptyl and the like. The term "halogen", "halo",


"halide" or "X" refers to a chlorine, bromine, or iodine atom and the term "Pg" refers to protecting group.

As used herein, the term "Ar-Y-" refers to a radical wherein Ar is an aryl group and Y is a C<sub>0</sub>-C<sub>4</sub> alkyl. The term "Ar" refers to a phenyl or naphthyl group unsubstituted or substituted with from one to three substituents selected from the group consisting of methylenedioxy, hydroxy, C<sub>1</sub>-C<sub>4</sub> alkoxy, fluoro and chloro. The term "C<sub>0</sub>-C<sub>4</sub> alkyl" refers to a saturated straight or branched chain hydrocarbyl radical of zero to four carbon atoms and includes a bond, methyl, ethyl, propyl, isopropyl, n-butyl, isobutyl, tertiary butyl and the like. Specifically included within the scope of the term "Ar-Y-" are phenyl, naphthyl, phenylmethyl or benzyl, phenylethyl, p-methoxybenzyl, p-fluorobenzyl and p-chlorobenzyl.

As used herein, the designation "~~" refers to a bond to a chiral atom for which the stereochemistry is not designated and compounds of Formula I wherein A is a bond is understood to be a 5-membered ring.

The compounds of Formula I wherein A is a bond, methylene, oxygen, sulfur or NH can be prepared by utilizing procedures and techniques well known and appreciated by one of ordinary skill in the art. A general synthetic scheme for preparing these compounds is set forth in Scheme A wherein all substituents, unless otherwise indicated, are previously defined.

Scheme A



A' = a bond, -CH<sub>2</sub>-, O, S or NH

The compounds of Formula I wherein A is a bond, methylene, oxygen, sulfur or NH can be prepared by reacting the appropriate protected thiol/carboxylic acid compound of structure 2 with the appropriate amino compound of structure 1. For example, the appropriate amino compound of structure 1 can be reacted with the appropriate protected thiol/carboxylic acid compound of structure 2 in the presence of a coupling reagent such as EDC (1-ethoxycarbonyl-2-ethoxy-1,2-dihydroquinoline), DCC (1,3-dicyclohexylcarbodiimide), or diethylcyanophosphonate in a suitable aprotic solvent, such as methylene chloride to give the appropriate tricyclic compound of structure 3.

Alternatively, the protected thiol/carboxylic acid compound of structure 2 can be converted to the corresponding, acid chloride, followed by reaction with the appropriate amino compound of structure 1 to give the appropriate compound of Formula 1.

The selection and utilization of suitable thiol protecting groups, such as t-butyl and 4-methoxybenzyl, is well known to one of ordinary skill in the art and are described in "Protective Groups in Organic Syntheses", Theodora W. Greene, Wiley (1981).

As summarized in Table 1, the R<sub>1</sub> and R<sub>2</sub> groups on the compounds Formula 1 can be manipulated using techniques and procedures well known and appreciated by one of ordinary skill in the art to give the corresponding compounds of structures 4 through 8.

For example, both the diphenylmethyl ester functionality and the thiol protecting group of the appropriate compound of structure 3 can be removed to give the appropriate carboxylic acid/thiol compound of structure 4.

For example, the appropriate tricyclic compound of structure 3 is contacted with a molar equivalent of mercuric acetate. The reactants are typically contacted in an appropriate acidic solvent such as trifluoroacetic acid. The reactants are typically stirred together at room temperature for a period of time ranging from 1-24 hours. Mercury is removed from the reaction mixture by the addition of excess hydrogen sulfide. The carboxylic acid/thiol compound of structure 4 is recovered from the reaction zone by extractive methods as is known in the art. It can be purified by silica gel chromatography.

The thiol functionality of the appropriate carboxylic acid/thiol compound of structure 4 can be alkylated to give the carboxylic acid/pivaloyloxymethyl thioacetal compound of structure 5.

For example, a carboxylic acid/pivaloyloxymethyl thioacetal compound of structure 5 can be prepared by treating the appropriate carboxylic acid/thiol compound of structure 4 with chloromethyl pivalate in a suitable aprotic solvent, such as dimethylformamide along with a non-nucleophilic base, such as cesium carbonate.

The thiol functionality of the appropriate carboxylic acid/thiol compound of structure 4 can be acylated to give the carboxylic acid/thioacetate or thiobenzoate compound of structure 6.

For example, the appropriate carboxylic acid/thiol compound of structure 4 can be contacted with a molar equivalent of an appropriate acylating agent such as acetic anhydride and a catalytic amount of an acid such as sulfuric acid. The reactants are typically stirred together for a period of time ranging from 10 minutes to 10 hours. The carboxylic acid/thioacetate compound of structure 6 can be recovered from the reaction zone by evaporation of the solvent. It can be purified by silica gel chromatography.

Alternatively, the appropriate carboxylic acid/thiol compound of structure 4 can be contacted with a molar equivalent of an appropriate acylating agent such as benzoyl chloride and a molar equivalent of a base such as pyridine. The reactants are typically stirred together for a period of time ranging from 10 minutes to 10 hours. The carboxylic acid/thiobenzoate compound of structure 6 can be recovered from the reaction zone by evaporation of the solvent. It can be purified by silica gel chromatography.

The carboxylic acid functionality of the appropriate carboxylic acid/thioacetate or thiobenzoate compound of structure 6 can be converted to the corresponding pivaloyloxymethyl ester/thioacetate or benzoate of structure 7 using techniques and procedures well known and appreciated in the art.

For example, a pivaloyloxymethyl ester/thioacetate or benzoate compound of structure 7 can be prepared by treating the carboxylic acid/thioacetate or thiobenzoate compound of structure 6 with 1 molar equivalent of chloromethyl pivalate in a suitable aprotic solvent, such as dimethylformamide along with a non-nucleophilic base, such as cesium carbonate.

The thioacetate or thiobenzoate functionality of the appropriate pivaloyloxymethyl ester/thioacetate or benzoate compound of structure 7 can be hydrolyzed to the corresponding pivaloyloxymethyl ester/thiol compounds of structure 8 with ammonia in a suitable protic solvent, such as methanol.

The appropriate carboxylic acid/thiol compound of structure 4 can be converted to the corresponding pivaloyloxymethyl ester/pivaloyloxymethyl thioacetal compound of structure 9.


For example, a pivaloyloxymethyl ester/pivaloyloxymethyl thioacetal compound of structure 9 can be prepared by treating the carboxylic acid/thiol compound of structure 4 with the 2 molar equivalents of chloromethyl pivalate in a suitable aprotic solvent, such as dimethylformamide along with a non-nucleophilic base, such as cesium carbonate.

TABLE 1

MANIPULATION OF R <sub>1</sub> AND R <sub>3</sub>		
Compound	R <sub>1</sub>	R <sub>3</sub>
<u>4</u>	H	H
<u>5</u>	CH <sub>2</sub> OCOC(CH <sub>3</sub> ) <sub>3</sub>	H
<u>6</u>	COCH <sub>3</sub> or COPh	H
<u>7</u>	COCH <sub>3</sub> or COPh	CH <sub>2</sub> OCOC(CH <sub>3</sub> ) <sub>3</sub>
<u>8</u>	H	CH <sub>2</sub> OCOC(CH <sub>3</sub> ) <sub>3</sub>
<u>9</u>	CH <sub>2</sub> OCOC(CH <sub>3</sub> ) <sub>3</sub>	CH <sub>2</sub> OCOC(CH <sub>3</sub> ) <sub>3</sub>

The compounds of Formula I wherein A is -NR<sub>4</sub> can be prepared by techniques and procedures well known and appreciated by one of ordinary skill in the art. A general synthetic procedure for preparing these compounds is set forth in Scheme B. In Scheme B, all substituents unless otherwise indicated are as previously defined.

Scheme B




Scheme B provides a general synthetic procedure for preparing the compounds of Formula I wherein A is -NR<sub>4</sub>. The amino functionality of the appropriate amino compound of structure 10 is subjected to

reductive alkylation with the appropriate aldehyde of structure 11 using sodium cyanoborohydride, as is well known in the art, to give the corresponding N-alkylamino compound of structure 12.

The  $R_1$  and  $R_3$  groups on the compounds Formula 1 wherein A is  $-NR_1$  can be manipulated as described previously in Scheme A and Table 1.

The compounds of Formula I wherein A is  $-NCOR_5$  can be prepared by techniques and procedures well known and appreciated by one of ordinary skill in the art. A general synthetic procedure for preparing these compounds is set forth in Scheme C. In Scheme C, all substituents unless otherwise indicated are as previously defined.

## Scheme C




Scheme C provides a general synthetic procedure for preparing the compounds of Formula I wherein A is  $-NCOR_5$ . The appropriate amino compound of structure 10 is acylated using the appropriate acyl chloride of structure 13 or the appropriate anhydride of structure 14, as is well known in the art, to give the corresponding N-acylamino compound of structure 15.

The groups  $R_1$  and  $R_3$  may be manipulated by techniques and procedures well known and appreciated in the art and described previously in Scheme A and shown in Table 1.

Starting materials for use in Scheme A through Scheme C are readily available to one of ordinary skill in the art. For example, certain tricyclic amino compounds of structure 1 wherein X is S are described in European Patent 0 249 223 (December 16, 1987) and certain other tricyclic amino compounds of structure 1 wherein A is methylene may be prepared as described in European Patent Application of Flynn and Beight, Application #34533AEP (June 11, 1987).


Tricyclic amino compounds of structure 1 wherein A is O may be prepared as described in Scheme D. In Scheme D, all substituents unless otherwise indicated are as previously defined.

## Scheme D





## Scheme D



Scheme D provides a general synthetic procedure for preparing amino compounds of structure 1 wherein A is O.

In step a, the appropriate phthalimide blocked (S)-phenylalanine derivative of structure 16 is converted to the corresponding acid chloride, then reacted with the appropriate L-serine methyl ester of structure 17 to give the corresponding 1-oxo-3-phenylpropyl-L-serine methyl ester of structure 18.

For example, the appropriate phthalimide blocked (S)-phenylalanine derivative of structure 16 can be reacted with oxalyl chloride in a suitable aprotic solvent, such as methylene chloride. The resulting acid chloride can then be coupled with the appropriate L-serine methyl ester of structure 17 using N-methylmorpholine in a suitable aprotic solvent, such as dimethylformamide, to give the appropriate 1-oxo-3-phenylpropyl-L-serine methyl ester of structure 18.

In step b, the hydroxy functionality of the appropriate 1-oxo-3-phenylpropyl-L-serine methyl ester of structure 18 is allylated with the allyl imidate of structure 19 to give the corresponding 1-oxo-3-phenylpropyl-L-serine-O-allyl methyl ester of structure 20.

For example, the appropriate 1-oxo-3-phenylpropyl-L-serine methyl ester of structure 18 is contacted with 2 molar equivalents of the allyl imidate of structure 19 and a molar equivalent of a suitable acid such as trifluoromethanesulfonic acid. The reactants are typically contacted in a suitable organic solvent mixture such as methylene chloride/cyclohexane. The reactants are typically stirred together at room temperature under an inert atmosphere for a period of time ranging from 2-24 hours. The 1-oxo-3-phenylpropyl-L-serine-O-allyl methyl ester of structure 20 is recovered from the reaction zone by extractive methods as is known in the art. It may be purified by silica gel chromatography or crystallization.

In step c, the appropriate 1-oxo-3-phenylpropyl-L-serine-O-allyl methyl ester of structure 20 is cyclized to give the corresponding (4S)-enamine of structure 21.

For example, the appropriate 1-oxo-3-phenylpropyl-L-serine-O-allyl methyl ester of structure 20 is first contacted with a molar excess of a mixture of ozone/oxygen. The reactants are typically contacted in a suitable organic solvent mixture such as methylene chloride/methanol. The reactants are typically stirred together for a period of time ranging from 5 minutes to 30 minutes or until a blue color persists and at a temperature range of from -78°C to -40°C. The reaction is quenched with an excess of methylsulfide and the intermediate aldehyde compound recovered from the reaction zone by extractive methods as is known in the art.

The intermediate aldehyde compound is then contacted with trifluoroacetic acid in a suitable aprotic solvent, such as methylene chloride to give the corresponding (4S)-enamine of structure 21.

In step d, the appropriate (4S)-enamine of structure 21 is cyclized to give the corresponding (4S)-tricyclic compound of structure 22 by an acid catalyzed Friedel-Crafts reaction. For example, the appropriate (4S)-enamine of structure 21 can be converted to the corresponding (4S)-tricyclic compound of structure 22 by treatment with a mixture of trifluoromethane sulfonic acid and trifluoroacetic anhydride in a suitable aprotic solvent, such as methylene chloride.


In step e, it may be necessary to reesterify the carboxy functionality due to the conditions of the work-up. For example, treatment of the crude product with bromodiphenylmethane in a suitable aprotic solvent,

such as dimethylformamide along with a non-nucleophilic base, such as cesium carbonate, may be used to give the corresponding (4S)-diphenylmethyl ester.


In step e, the phthalimide protecting group of the appropriate (4S)-tricyclic compound of structure 22 is removed to give the corresponding amino compound of structure 23 wherein X is O. For example, the phthalimide protecting group of the appropriate (4S)-tricyclic compound of structure 22 can be removed using hydrazine monohydrate in a suitable protic solvent such as methanol, to give the corresponding (4S)-tricyclic amino compound of structure 23.

Tricyclic amino compounds of structure 1 wherein A is NH may be prepared as described in Scheme E. In Scheme E, all substituents unless otherwise indicated are as previously defined.

Scheme E



## Scheme E



Scheme E provides an alternative general synthetic procedure for preparing amino compounds of structure 1 wherein A is NH.

In step a, the appropriate phthalimide blocked (S)-phenylalanine derivative of structure 16 is converted to the corresponding acid chloride, then reacted with the appropriate 3-trifluoroacetyl-amino-3-allyl-L-2-aminopropionic acid, methyl ester of structure 24 to give the corresponding 1-oxo-3-phenylpropyl-N-trifluoroacetyl-N-allyl-L-amino acid, methyl ester of structure 25 as described previously in Scheme D, step a.


In step b, the appropriate 1-oxo-3-phenylpropyl-N-trifluoroacetyl-N-allyl-L-amino acid methyl ester of structure 25 is cyclized to give the corresponding enamine of structure 26 as described previously in Scheme D, step c.

In step c, the appropriate (4S)-enamine of structure 26 is cyclized to give the corresponding (4S)-tricyclic compound of structure 27 as described previously in Scheme D, step d.


In step d, the phthalimide protecting group of the appropriate (4S)-tricyclic compound of structure 27 is removed to give the corresponding (4S)-amino compound of structure 28 wherein X is NH as described in Scheme D, step e.

Tricyclic amino compounds of structure 1 wherein A is methylene may be prepared as described in Scheme F. In Scheme F, all substituents unless otherwise indicated are as previously defined.

## Scheme F



## Scheme F cont.



Scheme F provides a general synthetic procedure for preparing the tricyclic amino compounds of structure 1 wherein A is methylene.

In step a, the appropriate phthalimide blocked (S)-phenylalanine derivative of structure 16 can be converted to the corresponding acid chloride, then reacted with the appropriate amino acid methyl ester of structure 29 in a coupling reaction. For example, the appropriate phthalimide blocked (S)-phenylalanine derivative of structure 16 can be reacted with oxalyl chloride in a suitable aprotic solvent, such as methylene chloride. The resulting acid chloride can then be coupled with the appropriate amino acid methyl ester of structure 29 using N-methylmorpholine in a suitable aprotic solvent, such as dimethylformamide, to give the appropriate 1-oxo-3-phenylpropyl-amino acid methyl ester derivative of structure 30.

In step b, the hydroxymethylene functionality of the appropriate 1-oxo-3-phenylpropyl-amino acid methyl ester derivative of structure 30 can be oxidized to the appropriate aldehyde of structure 31 by oxidation techniques well known and appreciated in the art. For example, the hydroxymethylene functionality of the appropriate 1-oxo-3-phenylpropyl-amino acid methyl ester derivative of structure 30 can be oxidized to the appropriate aldehyde of structure 31 by means of a Swern oxidation using oxalyl chloride and dimethylsulfoxide in a suitable aprotic solvent, such as methylene chloride.

In step c, the appropriate aldehyde of structure 31 can be cyclized to the appropriate enamine of structure 32 by acid catalysis. For example, the appropriate aldehyde of structure 31 can be cyclized to the appropriate enamine of structure 32 by treatment with trifluoroacetic acid in a suitable aprotic solvent, such as methylene chloride.


In step d, the appropriate enamine of structure 32 can be converted to the corresponding tricyclic compound of structure 33 by an acid catalyzed Friedel-Crafts reaction. For example, the appropriate enamine of structure 32 can be converted to the corresponding tricyclic compound of structure 33 by treatment with a mixture of trifluoromethane sulfonic acid and trifluoroacetic anhydride in a suitable aprotic solvent, such as methylene chloride.

In step d, it may be necessary to reesterify the carboxy functionality due to the conditions of the work-up. For example, treatment of the crude product with bromodiphenylmethane in a suitable aprotic solvent, such as dimethylformamide along with a non-nucleophilic base, such as cesium carbonate, may be used to give the corresponding diphenylmethyl ester.

In step e, the phthalimide protecting group of the appropriate tricyclic compound of structure 33 can be removed using techniques and procedures well known in the art. For example, the phthalimide protecting group of the appropriate tricyclic compound of structure 33 can be removed using hydrazine monohydrate in a suitable protic solvent such as methanol, to give the corresponding amino compound of structure 34.

The compounds of Formula I wherein A is a bond can be prepared by utilizing procedures and techniques well known and appreciated by one of ordinary skill in the art. A general synthetic scheme for preparing these compounds is set forth in Scheme G wherein all substituents, unless otherwise indicated, are previously defined.

Scheme G



Scheme G provides a general synthetic procedure for preparing compounds of Formula I wherein A is a bond.

In step a, the N-(phenylmethylene)glycine methyl ester of structure 35 can be treated with one equivalent of a non-nucleophilic base, such as lithium diisopropylamide, in a suitable aprotic solvent, such as tetrahydrofuran, followed by addition of a 4-halobutene of structure 36 to give 2-(3-butenyl)-N-(phenylmethylene)glycine methyl ester of structure 37.

In step b, the N-(phenylmethylene) functionality of 2-(3-butenyl)-N-(phenylmethylene)glycine methyl ester of structure 37 can be hydrolyzed under acidic conditions, such as with hydrochloric acid in a suitable aprotic solvent, such as ethyl ether to give 2-(3-butenyl)-glycine methyl ester of structure 38.

In step c, the appropriate amide compound of structure 38 can be prepared by reacting the appropriate phthalimide protected (S)-phenylalanine compound of structure 16 with 2-(3-butenyl)-glycine methyl ester of

structure 39 under coupling reaction conditions, such as with EEDQ, in a suitable aprotic solvent, such as methylene chloride.

In step d, the olefin functionality of the appropriate amide compound of structure 39 can be converted to the appropriate aldehyde compound of structure 40 under conditions of oxidative cleavage, such as treatment with ozone in a suitable solvent mixture, such as methylene chloride and methanol.

The compounds of Formula I wherein A is a bond can be prepared from an appropriate aldehyde of structure 40 in a process as outlined previously in Scheme F, steps c-e and Scheme A.

The individual 3(S) and 3(R) esters of the compounds of Formula I wherein A is a bond, R<sub>1</sub> is t-butyl and R<sub>2</sub> is methyl can be prepared from an appropriate aldehyde of structure 40 in a process as outlined previously in Scheme F, step c, separating the 3(S) and 3(R) esters of the enamine compounds formed from the cyclization reaction described in Scheme F, step c and completing the process as outlined in Scheme F, steps d-e and Scheme A.

The groups R<sub>1</sub> and R<sub>2</sub> may be manipulated by techniques and procedures well known and appreciated in the art and described previously in Scheme A and Table 1.

Starting materials for use in the general synthetic procedures outlined in Schemes D and G are readily available to one of ordinary skill in the art. For example, N<sup>o</sup>-(benzyloxycarbonyl)-β-(amino)-L-alanine is described in *J. Am.Chem.Soc.*, 107(24) 7105 1985, t-butyl chloromethylsulfide is described in *J.Am.Chem.Soc.* 77 572 1955 and allyl trichloroacetimidate is described in *J.Chem.Soc.Perkin Trans.* 1(11) 2247 1985. 3,4-(Dicarboethoxy)thiophene is described in *Bull.Chem.Soc.Jpn.* 41 2532 1968, N-(phenylmethylene)glycine methyl ester is described in *J.Org.Chem.* 41, 3491 1976 and 3,4-(dicarboethoxy)-pyrrole is described in *Tetrahedron Lett.* 3165 1971.

The following examples present typical syntheses as described in Schemes A through G. These examples are understood to be illustrative only and are not intended to limit the scope of the present invention in any way. As used herein, the following terms have the indicated meanings: "g" refers to grams; "mmol" refers to millimoles; "mL" refers to milliliters; "bp" refers to boiling point; "°C" refers to degrees Celsius; "mm Hg" refers to millimeters of mercury; "μL" refers to microliters; "μg" refers to micrograms; and "μM" refers to micromolar.

#### Example 1

[4S-[4α, 7α(R), 12bβ]]-7-[(2-Thio-2-oxoindan)methylamino]-1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido[2,1-a]-[2]benzazepine-4-carboxylic acid

Scheme F, Step a: (S)-N-[2-(1,3-Dihydro-1,3-dioxo-2H-isoindol-2-yl)-1-oxo-3-phenylpropyl]-6-hydroxy-(S)-norleucine, methyl ester

Mix phthalic anhydride (1.82kgs, 12.3mole), (S)-phenylalanine (1.84kgs, 11.1 moles) and anhydrous dimethylformamide (2.26L). Stir at 115-120°C for 2 hours under a nitrogen atmosphere. Pour into rapidly stirring water (32.6L) and cool overnight at 0°C. Filter, wash with cold water (2X2L) and air dry. Dissolve in a mixture of 9A ethanol (8.05L) and water (8.05L) and heat at reflux temperature. Gravity filter, cool to ambient temperature and refrigerate overnight at about 0°C. Filter the crystallized product, wash with cold 50:50 9A ethanol/water (2X2L) and air dry to yield 2.96kg (90.3%) of N-phthaloyl-(S)-phenylalanine; mp 177-179°C.

Mix N-phthaloyl-(S)-phenylalanine (50.2g, 0.17mole), methylene chloride (660mL) and dimethylformamide (0.5mL) under a nitrogen atmosphere. Add oxalyl chloride (17.7mL, 0.2mole) over about 5 minutes with stirring. Stir at ambient temperature for 3 hours and evaporate the solvent in vacuo to leave N-phthaloyl-(S)-phenylalanine, acid chloride as a solid (54.3g, 101.9%).

Mix 6-hydroxy-(S)-norleucine, methyl ester, hydrochloride salt (33.5g, 0.1mole) and dimethylformamide (201mL), cool to about 0°C and place under nitrogen atmosphere. Add by dropwise addition, N-methylmorpholine (51mL, 0.46mole) with cooling so that the pot temperature stays at 0-5°C. Stir at 0-5°C for an additional 10 minutes, then add a solution of N-phthaloyl-(S)-phenylalanine, acid chloride (53.5g, 0.17mole) in methylene chloride (270mL) over 30 minutes with cooling so that the temperature stays at 0-5°C. Remove the cooling bath and stir at room temperature for 18 hours.

Evaporate the methylene chloride in vacuo and dilute the remaining residue with ethyl acetate (800mL). Extract the resulting mixture with water (800mL), separate the organic layer and extract with 1N hydrochloric acid (270mL), followed by water (3X500mL). Dry the organic layer (MgSO<sub>4</sub>), filter and evaporate in vacuo to yield crude product (76g, 98%). Dissolve the crude product in hot toluene (223.5mL), cool to room temperature, then cool overnight at about 0°C. Filter the crystallized product, wash with cold toluene and air

dry to yield 56.6g (76%) of the title compound; mp 128-130 °C.

Scheme F, Step b: 2-(1,3-Dihydro-1,3-dioxo-2H-isoindol-2-yl)-1-oxo-3-phenylpropyl-6-oxo-(S)-norleucine, methyl ester

5 Mix oxalyl chloride (80mL, 0.92mole) and methylene chloride (2L) and place under nitrogen atmosphere. Cool below -50 °C and add a solution of dimethyl sulfoxide (65.4mL, 0.92mole) in methylene chloride (425mL). Stir for 15 minutes and add a solution of (S)-N-[2-(1,3-dihydro-1,3-dioxo-2H-isoindol-2-yl)-1-oxo-3-phenylpropyl]-6-hydroxy-(S)-norleucine, methyl ester (200g, 0.456mole) in methylene chloride (800mL) over  
10 about 45 minutes, keeping the pot temperature below -50 °C for 30 minutes. Add triethylamine (420mL, 3.01mole) over 30 minutes. Stir while warming to 0 °C over 1.25 hours. Transfer the reaction mixture to a 12-liter flask. Stir and cool while adding a solution of OXONE (potassium peroxymonosulfate) (566g) in water (6.74L) at such a rate that the pot temperature stays below 15 °C. Stir for 5 minutes, separate the organic layer and extract the aqueous layer with methylene chloride (1L). Combine the organic phases, dry  
15 (MgSO<sub>4</sub>) and filter to yield the title compound as a solution.

Scheme F, Step c: [S-(R\*,R\*)]-N-[2-(1,3-Dihydro-1,3-dioxo-2H-isoindol-2-yl)-1-oxo-3-phenylpropyl]-1,2,3,4-tetrahydro-2-pyridinecarboxylic acid, methyl ester

20 Transfer the solution of 2-(1,3-dihydro-1,3-dioxo-2H-isoindol-2-yl)-1-oxo-3-phenylpropyl-6-oxo-(S)-norleucine, methyl ester in methylene chloride (volume about 4.5L) to a 12-liter flask and place under nitrogen atmosphere. Stir and add trifluoroacetic acid (440mL, 5.71mole) in one portion. Stir the resulting mixture at room temperature for one hour, then rapidly cool to about 0 °C. Add a solution of sodium hydroxide (240g, 6.0mole) in water (3.4L) in a slow stream to the vigorously stirred mixture at such a rate that the pot  
25 temperature stays at about 0 °C. Separate the organic phase and extract the aqueous phase with methylene chloride (1L). Combine the organic phases and dry (MgSO<sub>4</sub>). Filter and remove the solvent in vacuo to leave a residue (262g, 137%).

Dissolve the above residue in diethyl ether (1L) and wash with water (5X500mL). Evaporate the organic phase in vacuo to leave a residue of 229g. Dilute the residue with methylene chloride (200mL) and purify by  
30 silica gel chromatography (methylene chloride) to yield a viscous residue of 225g.

Dilute the above residue with diethyl ether (250mL) and allow to stand at room temperature for 24 hours. Filter the solid, wash with diethyl ether, and air dry to yield 123.2g; mp 140-142.5 °C. Recrystallize (methylene chloride (125mL)/isopropanol (615mL)) by boiling off the solvent until the pot temperature reaches 75 °C and allowing the resulting sample to stand at room temperature for 24 hours. Filter, wash  
35 with cold isopropanol and air dry to yield 101.5g of the title compound; mp 144-146 °C.

Evaporate the filtrate from the 101.5g in vacuo to yield 24g. Recrystallize (isopropanol) to yield an additional 3.5g of the title compound.

Evaporate the filtrate from the 123.2g in vacuo to leave 62g of oil. Purify by silica gel chromatography (25% ethyl acetate/75% hexane), collecting 21-500mL fractions. Combine fractions 9-20 and evaporate in vacuo  
40 to yield 35g of a viscous oil. Recrystallize three times (isopropanol/5mL/g) to yield an additional 11.9g of the title compound; mp 142.5-144.5 °C. Total yield of useful material: 116.9g (61.3%).

Scheme F, Step d: [4S-[4 $\alpha$ , 7 $\alpha$ (R\*), 12b $\beta$ ]]-7-[(1,3-Dihydro-1,3-dioxo-2H-isoindol-2-yl)]-1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido[2,1-a][2]benzazepine-4-carboxylic acid, diphenylmethyl ester

45 Mix trifluoromethanesulfonic acid (500g, 3.33mole) and trifluoroacetic anhydride (74.8mL, 0.53mole) and place under nitrogen atmosphere. Stir and add a solution of [S-(R\*,R\*)]-N-[2-(1,3-dihydro-1,3-dioxo-2H-isoindol-2-yl)-1-oxo-3-phenylpropyl]-1,2,3,4-tetrahydro-2-pyridinecarboxylic acid, methyl ester (200g, 0.48mole) in methylene chloride (1L) with cooling at such a rate as to keep the pot temperature below  
50 35 °C. Stir at ambient temperature for 2 days. Pour into vigorously stirring ice water (5L) and stir for 30 minutes. Extract with ethyl acetate (3X1L), combine the organic phases and wash with water (3x500mL). Evaporate in vacuo to a residue. Dissolve the residue in ethyl acetate (4L) and extract with 1/4 saturated potassium hydrogen carbonate (1L), then 1/3 saturated potassium hydrogen carbonate (7X1L). Combine the aqueous extracts and dilute with ethyl acetate (2L). Stir the resulting mixture and cool to 5-10 °C. Adjust to  
55 pH 2 using concentrated hydrochloric acid (about 750mL).

Separate the organic phase and extract the aqueous phase with ethyl acetate (3X1L). Combine the ethyl acetate extracts, wash with water (3X1L), then saturated sodium chloride (0.8L), and dry (MgSO<sub>4</sub>). Filter and wash with ethyl acetate (3X200mL). Evaporate in vacuo to leave (188.3g, 101.5%) [4S-[4 $\alpha$ , 7 $\alpha$ (R\*), 12b $\beta$ ]]-7-



((1,3-dihydro-1,3-dioxo-2H-isoindol-2-yl))-1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido[2,1-a][2]benzazepino-4-carboxylic acid as a colorless foam.

Dissolve [4S-[4 $\alpha$ , 7 $\alpha$ (R\*), 12b $\beta$ ]]-7-[(1,3-dihydro-1,3-dioxo-2H-isoindol-2-yl))-1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido[2,1-a][2]benzazepine-4-carboxylic acid (113.9g, 0.28mole) in methylene chloride (1.2L) and dry over anhydrous MgSO<sub>4</sub> (60g). Filter and wash with methylene chloride (3X200mL). Evaporate in vacuo to a residue. Dissolve the residue in anhydrous dimethylformamide (860mL) and place under nitrogen atmosphere. Add cesium carbonate (98.9g, 0.3mole) in one portion. Stir for 45 minutes at ambient temperature. Add bromodiphenylmethane (164.8g, 0.67mole). Stir the resulting mixture at ambient temperature for 18 hours. Quench the reaction with ethyl acetate (2.464L) and water (630mL). Separate the organic phase and wash with water (7X625mL), 1/4 saturated potassium hydrogen carbonate (625mL), water (625mL), and saturated sodium chloride (625mL). Dry (MgSO<sub>4</sub>), filter and evaporate in vacuo to yield 214.4g of an oil. Extract the combined aqueous washings with ethyl acetate (3X500mL), wash with water (4X300mL) and dry (MgSO<sub>4</sub>). Filter and evaporate in vacuo to yield an additional 20.2g of an oil. Dissolve the crude product (234.6g) in methylene chloride (200mL) and plug filter through 213g of silica gel, eluting with methylene chloride (2L). Boil off the solvent and replace with hexane (3L), with the pot temperature reaching a maximum of 65°C. Cool to ambient temperature, decant off the precipitated oil and crystallize (9A ethanol) to yield 96.6g (60%) of the title compound; mp 153-155°C.

Scheme F, Step e: [4S-[4 $\alpha$ , 7 $\alpha$ (R\*), 12b $\beta$ ]]-7-(Amino)-1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido[2,1-a][2]benzazepine-4-carboxylic acid, diphenylmethyl ester

Mix [4S-[4 $\alpha$ , 7 $\alpha$ (R\*), 12b $\beta$ ]]-7-[(1,3-dihydro-1,3-dioxo-2H-isoindol-2-yl))-1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido[2,1-a][2]benzazepine-4-carboxylic acid, diphenylmethyl ester (170.9g, 0.3mole), hydrazine monohydrate (34.4g, 0.68mole) and methanol (3.4L) under nitrogen atmosphere. Heat at reflux for 5 hours. Cool to ambient temperature and filter to remove phthaloyl hydrazide. Evaporate the filtrate in vacuo to a residue and slurry in chloroform (600mL). Remove insoluble phthaloyl hydrazide by filtration and wash with chloroform (4X210mL). Wash the filtrate with water (4X429mL), dry (MgSO<sub>4</sub>), and filter. Evaporate the filtrate to a solid residue of the title compound weighing 142g (107.7%).

Scheme A: [4S-[4 $\alpha$ , 7 $\alpha$ (R\*), 12b $\beta$ ]]-7-[(2-Thio-2-oxoindan)methylamino]-1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido[2,1-a][2]benzazepine-4-carboxylic acid

Dissolve diethylmalonate (15.2mL, 0.100mol) in tetrahydrofuran (800mL). Cool in an ice bath and treat with sodium hydride (3.0g, 0.10mol, 80% in mineral oil). Stir until solution attained and add  $\alpha,\alpha'$ -dibromo-o-xylene (26.4g, 0.100mol). Stir for 30 minutes then add additional sodium hydride (3.0g, 0.10mol). Stir at room temperature for 20 hours, filter through filter aid and evaporate the solvent in vacuo. Purify by silica gel chromatography to give 2,2-(dicarboethoxy)indan as a pale yellow oil (16.0g, 61.5%).

Dissolve 2,2-(dicarboethoxy)indan (15.9g, 60.6mmol) in dimethylsulfoxide (140mL). Add water (14mL) and lithium chloride (7.0g, 0.16mol). Heat at reflux for 4 hours, cool and partition between water (150mL) and methylene chloride (2X150mL). Wash the organic phase with water (150mL), dry (MgSO<sub>4</sub>) and pass through a silica gel plug to give 2-(carboethoxy)indan as an amber oil (6.49g, 56%).

Dissolve 2-(carboethoxy)indan (6.49g, 34.1mmol) in ethanol (95%, 150mL) and water (75mL). Add potassium hydroxide (9.5g, 0.17mol) and stir at room temperature for 1 hour. Partition between water (150mL) and ethyl ether (2X150mL). Acidify the aqueous phase with hydrochloric acid to pH 1. Extract with methylene chloride (2X150mL), dry (Na<sub>2</sub>SO<sub>4</sub>) and evaporate the solvent in vacuo to give 2-indancarboxylic acid as a tan solid (3.82g, 69%).

Dissolve 2-indancarboxylic acid (3.82g, 23.5mmol) in methanol (60mL) and treat with dimethoxypropane (5.8mL, 47mmol) and sulfuric acid (0.8mL). Stir at room temperature for 6 days. Evaporate the solvent in vacuo, dilute with methylene chloride (75mL) and wash with saturated sodium hydrogen carbonate (35mL). Extract the aqueous phase with methylene chloride (30mL), wash combined organics with brine (30mL) and dry (Na<sub>2</sub>SO<sub>4</sub>). Evaporate the solvent in vacuo and pass through a plug of silica gel to give 2-(carbomethoxy)indan as a yellow oil (3.98g).

Mix 4-methoxybenzylthiol (3.0g, 19mmol) in sodium hydroxide (20mL of a 2.5N aqueous solution) and methanol (10mL). Add saturated aqueous copper sulfate solution (1.5mL) and stir at room temperature for 2 hours, blowing air over the top of the mixture. Filter the solid, wash with water and dry to give 4-methoxybenzyl disulfide as a pale yellow powder (2.71g, 91%).

Cool lithium hexamethyldisilazane (4.2mL, 4.2mol, 1.0M in tetrahydrofuran) to -78°C and treat with a solution of 2-(carbomethoxy)indan (625mg, 3.55mmol) in tetrahydrofuran (5mL). Stir for 1 hour add

hexamethylphosphoramide (0.93mL, 5.3mmol) and stir for 5 minutes. Add 4-methoxybenzyl disulfide (1.6g, 5.2mmol) in tetrahydrofuran (10mL). Stir for 5 hours at -78 °C and quench with a solution of ammonium chloride. Partition between ethyl acetate (75mL) and brine (30mL). Dry (Na<sub>2</sub>SO<sub>4</sub>), evaporate the solvent in vacuo and purify by silica gel chromatography (4:1 hexane/ethyl acetate) to give 2-(carbomethoxy)-2-(4-methoxybenzylthio)indan (2.20g).

Dissolve 2-(carbomethoxy)-2-(4-methoxybenzylthio)indan (2.20g, 3.55mmol) in 95% ethanol (25mL), water (12mL) and tetrahydrofuran (15mL). Treat with potassium hydroxide (1.3g, 23mmol) and stir at room temperature for 1 hour. Filter and evaporate the solvent in vacuo. Partition between water (125mL) and ether (75mL). Separate the aqueous phase and acidify with cold concentrated hydrochloric acid. Extract with methylene chloride (75mL), dry (Na<sub>2</sub>SO<sub>4</sub>) and evaporate the solvent in vacuo. Purify by silica gel chromatography (2:1 hexane/ethyl acetate) to give 2-carboxy-2-(4-methoxybenzylthio)indan as a yellow solid (0.48g).

Dissolve [4S-[4 $\alpha$ , 7 $\alpha$ (R), 12b $\beta$ ]]-7-(amino)-1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido[2,1-a][2]benzazepine-4-carboxylic acid, diphenylmethyl ester (200mg, 0.454mmol) and EDC (130mg, 0.68mmol) in tetrahydrofuran (5mL). Treat with 2-carboxy-2-(4-methoxybenzylthio)indan (143mg, 0.454mmol). Stir at room temperature for 20 hours and evaporate the solvent in vacuo. Dissolve the residue in ethyl acetate (50mL) and wash with 5% sulfuric acid (15mL) then saturated sodium hydrogen carbonate (15mL). Dry (Na<sub>2</sub>SO<sub>4</sub>) and evaporate the solvent in vacuo. Purify by silica gel chromatography (2.5:1 hexane/ethyl acetate to 3:2 hexane/ethyl acetate) to give [4S-[4 $\alpha$ , 7 $\alpha$ (R), 12b $\beta$ ]]-7-[(2-(4-methoxybenzylthio)-2-oxoindan)methylamino]-1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido[2,1-a][2]benzazepine-4-carboxylic acid, diphenylmethyl ester as a pale yellow oil (245mg, 73.1%).

Dissolve [4S-[4 $\alpha$ , 7 $\alpha$ (R), 12b $\beta$ ]]-7-[(2-(4-methoxybenzylthio)-2-oxoindan)methylamino]-1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido[2,1-a][2]benzazepine-4-carboxylic acid, diphenylmethyl ester (120mg, 0.163mmol) in methylene chloride (3mL) and cool to 0 °C. Treat with trifluoroacetic acid (1.5mL), anisole (0.19mL, 1.7mmol), and mercuric acetate (65mg, 0.2mmol). Stir at 0 °C for 3 hours then bubble hydrogen sulfide gas through the solution for 10 minutes. Filter and wash with methylene chloride. Wash organic phase with water (20mL), dry (Na<sub>2</sub>SO<sub>4</sub>) and evaporate the solvent in vacuo. Purify by silica gel chromatography (2.5:1 hexane/ethyl acetate to 1:1 hexane/ethyl acetate with 1% acetic acid) to give the title compound (65mg, 89%).

#### Example 2

[4S-[4 $\alpha$ , 7 $\alpha$ (R), 12b $\beta$ ]]-7-[(2-Pivaloyloxymethylthio)-2-oxoindan)methylamino]-1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido[2,1-a][2]benzazepine-4-carboxylic acid

Dissolve [4S-[4 $\alpha$ , 7 $\alpha$ (R), 12b $\beta$ ]]-7-[(2-thiol-2-oxoindan)methylamino]-1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido[2,1-a][2]benzazepine-4-carboxylic acid (126mg, 0.28mmol) in methylene chloride (1mL) and dry over anhydrous MgSO<sub>4</sub> (60mg). Filter and wash with methylene chloride (3X20mL). Evaporate in vacuo to a residue. Dissolve the residue in anhydrous dimethylformamide (10mL) and place under nitrogen atmosphere. Add cesium carbonate (100mg, 0.3mmol) in one portion. Stir for 45 minutes at ambient temperature. Add chloromethyl pivalate (42g, 0.28mmol). Stir the resulting mixture at ambient temperature for 18 hours. Quench the reaction with ethyl acetate (3mL) and water (10mL). Separate the organic phase and wash with water (7X10mL), 1/4 saturated potassium hydrogen carbonate (10mL), water (10mL), and saturated sodium chloride (10mL). Dry (MgSO<sub>4</sub>), filter and evaporate in vacuo to yield the title compound.

#### Example 3

[4S-[4 $\alpha$ , 7 $\alpha$ (R), 12b $\beta$ ]]-7-[(2-Thioacetate)-2-oxoindan)methylamino]-1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido[2,1-a][2]benzazepine-4-carboxylic acid

Mix [4S-[4 $\alpha$ , 7 $\alpha$ (R), 12b $\beta$ ]]-7-[(2-thiol-2-oxoindan)methylamino]-1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido[2,1-a][2]benzazepine-4-carboxylic acid (2.73g, 6.07mmol) and sulfuric acid (0.1mL of a 10% solution in acetic acid). Add acetic anhydride (0.572mL, 6.07mmol) and stir for 2 hours. Dilute with ethyl ether, wash with water then saturated sodium hydrogen carbonate and brine. Dry (MgSO<sub>4</sub>), evaporate the solvent in vacuo and purify by silica gel chromatography to give the title compound.

## Example 4

[4S-[4 $\alpha$ , 7 $\alpha$ (R'), 12b $\beta$ ]]-7-[(2-Thioacetate-2-oxoindan)methylamino]-1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido[2,1-a][2]benzazepine-4-carboxylic acid, pivaloyloxymethyl ester

Dissolve [4S-[4 $\alpha$ , 7 $\alpha$ (R'), 12b $\beta$ ]]-7-[(2-thioacetate-2-oxoindan)methylamino]-1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido[2,1-a][2]benzazepine-4-carboxylic acid (138g, 0.28mmol) in methylene chloride (1mL) and dry over anhydrous MgSO<sub>4</sub> (60mg). Filter and wash with methylene chloride (3X20mL). Evaporate in vacuo to a residue. Dissolve the residue in anhydrous dimethylformamide (10mL) and place under nitrogen atmosphere. Add cesium carbonate (100mg, 0.3mmol) in one portion. Stir for 45 minutes at ambient temperature. Add chloromethyl pivalate (42g, 0.28mmol). Stir the resulting mixture at ambient temperature for 18 hours. Quench the reaction with ethyl acetate (3mL) and water (10mL). Separate the organic phase and wash with water (7X10mL), 1/4 saturated potassium hydrogen carbonate (10mL), water (10mL), and saturated sodium chloride (10mL). Dry (MgSO<sub>4</sub>), filter and evaporate in vacuo to yield the title compound.

## Example 5

[4S-[4 $\alpha$ , 7 $\alpha$ (R'), 12b $\beta$ ]]-7-[(2-Thio-2-oxoindan)methylamino]-1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido[2,1-a][2]benzazepine-4-carboxylic acid, pivaloyloxymethyl ester

Stir [4S-[4 $\alpha$ , 7 $\alpha$ (R'), 12b $\beta$ ]]-7-[(2-thioacetate-2-oxoindan)methylamino]-1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido[2,1-a][2]benzazepine-4-carboxylic acid, pivaloyloxymethyl ester (2.42g, 4mmol) and saturated methanolic ammonia at ambient temperature until hydrolysis is complete. Evaporate the solvent in vacuo and purify by silica gel chromatography to give the title compound.

## Example 6

[4S-[4 $\alpha$ , 7 $\alpha$ (R'), 12b $\beta$ ]]-7-[(2-Pivaloyloxymethylthio-2-oxoindan)methylamino]-1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido[2,1-a][2]benzazepine-4-carboxylic acid, pivaloyloxymethyl ester

Dissolve [4S-[4 $\alpha$ , 7 $\alpha$ (R'), 12b $\beta$ ]]-7-[(2-thiol-2-oxoindan)methylamino]-1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido[2,1-a][2]benzazepine-4-carboxylic acid (63mg, 0.14mole) in methylene chloride (1mL) and dry over anhydrous MgSO<sub>4</sub> (60mg). Filter and wash with methylene chloride (3X20mL). Evaporate in vacuo to a residue. Dissolve the residue in anhydrous dimethylformamide (10mL) and place under nitrogen atmosphere. Add cesium carbonate (100mg, 0.3mmol) in one portion. Stir for 45 minutes at ambient temperature. Add chloromethyl pivalate (42g, 0.3mmol). Stir the resulting mixture at ambient temperature for 18 hours. Quench the reaction with ethyl acetate (3mL) and water (10mL). Separate the organic phase and wash with water (7X10mL), 1/4 saturated potassium hydrogen carbonate (10mL), water (10mL), and saturated sodium chloride (10mL). Dry (MgSO<sub>4</sub>), filter and evaporate in vacuo to yield the title compound.

## Example 7

[4S-[4 $\alpha$ , 7 $\alpha$ (R'), 12b $\beta$ ]]-7-[(2-(2-mercaptoethyl)-2-oxoindan)methylamino]-1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido[2,1-a][2]benzazepine-4-carboxylic acid

Cool lithium hexamethyldisilazane (4.2mL 4.2mmol) in a dry ice/acetone bath and add, by dropwise addition, a solution of 2-(carbomethoxy)indane (625mg, 3.55mmol) in tetrahydrofuran (6mL). Stir for 30 minutes then add hexamethylphosphoramide (1.0mL, 5.7mmol). Stir for 45 minutes then add 1-bromo-2-chloroethane (0.37mL, 4.44mmol). Stir for 3 hours, remove the ice bath and allow to warm to room temperature. Quench with aqueous ammonium chloride (30mL) and extract with ethyl acetate (75mL). Dry (Na<sub>2</sub>SO<sub>4</sub>) and evaporate the solvent in vacuo. Pass through a plug of silica gel (6:1 hexane/ethyl acetate) to give 2-(carbomethoxy)-2-(2-chloroethane)indane (0.62g).

Dissolve 4-methoxybenzylthiol (0.7mL, 5mmol) in tetrahydrofuran (10mL) and place under an argon atmosphere. Treat with sodium hydride (144mg of an 80% dispersion in mineral oil, 4.8mmol). Stir the suspension for 10 minutes and add tetrabutylammonium iodide (40mg, 0.11mmol). Stir for 10 minutes then add a solution of 2-(carbomethoxy)-2-(2-chloroethane)indane (0.6g, 3.55mmol) in tetrahydrofuran (5mL). Stir for 18 hours and partition between aqueous ammonium chloride (15mL) and ethyl acetate (50mL). Dry (Na<sub>2</sub>SO<sub>4</sub>) and evaporate the solvent in vacuo. Purify by silica gel chromatography (6:1 hexane/ethyl acetate)

to give 2-(carbomethoxy)-2-[2-(4-methoxybenzylthio)ethane]indan (0.75g).

Dissolve 2-(carbomethoxy)-2-[2-(4-methoxybenzylthio)ethane]indan (0.75g, 3.55mmol) in 95% ethanol (16mL), water (8mL) and tetrahydrofuran (10mL). Treat with potassium hydroxide (1.3g, 23mmol). Stir for 1 hour and evaporate the solvent in vacuo. Partition between water (50mL) and ethyl ether (2X35mL). Cool the aqueous phase in an ice bath and acidify with concentrated hydrochloric acid to pH 1. Extract with methylene chloride (75mL), dry (Na<sub>2</sub>SO<sub>4</sub>) and evaporate the solvent in vacuo to give 2-carboxy-2-[2-(4-methoxybenzylthio)ethane]indan as a white solid (147mg).

Dissolve [4S-[4 $\alpha$ , 7 $\alpha$ (R\*), 12b $\beta$ ]]-7-(amino)-1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido[2,1-a][2]benzazepine-4-carboxylic acid, diphenylmethyl ester (180mg, 0.409mmol) and EDC (118mg, 0.813mmol) in tetrahydrofuran (5mL). Treat with a solution of 2-carboxy-2-[2-(4-methoxybenzylthio)ethane]indan (140mg, 0.409mmol) in tetrahydrofuran (3mL). Stir at room temperature for 20 hours and evaporate the solvent in vacuo. Dissolve the residue in ethyl acetate (40mL) and wash with 5% sulfuric acid (15mL) then saturated sodium hydrogen carbonate (15mL). Dry (Na<sub>2</sub>SO<sub>4</sub>) and evaporate the solvent in vacuo. Purify by silica gel chromatography (2:1 hexane/ethyl acetate to 3:2 hexane/ethyl acetate) to give [4S-[4 $\alpha$ , 7 $\alpha$ (R\*), 12b $\beta$ ]]-7-[(2-[2-(4-methoxybenzylthio)ethane-2-oxoindan)methylamino]-1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido[2,1-a][2]benzazepine-4-carboxylic acid, diphenylmethyl ester as a white solid (168mg, 53.2%).

Dissolve [4S-[4 $\alpha$ , 7 $\alpha$ (R\*), 12b $\beta$ ]]-7-[(2-[2-(4-methoxybenzylthio)ethane-2-oxoindan)methylamino]-1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido[2,1-a][2]benzazepine-4-carboxylic acid, diphenylmethyl ester (166mg, 0.217mmol) in methylene chloride (4mL) and cool to 0°C. Treat with trifluoroacetic acid (2mL), anisole (0.24mL, 2.2mmol), and mercuric acetate (86mg, 0.27mmol). Stir at 0°C for 3 hours then bubble hydrogen sulfide gas through the solution for 10 minutes. Filter and dilute with methylene chloride. Wash organic phase with water (20mL), dry (Na<sub>2</sub>SO<sub>4</sub>) and evaporate the solvent in vacuo. Purify by silica gel chromatography (2:1 hexane/ethyl acetate to 1:1 hexane/ethyl acetate with 1% acetic acid) to give the title compound (78mg, 75%).

#### Example 8

[4S-[4 $\alpha$ , 7 $\alpha$ (R\*), 12b $\beta$ ]]-7-[(2-Mercaptomethyl-2-oxoindan)methylamino]-1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido[2,1-a][2]benzazepine-4-carboxylic acid

Dissolve diethyl malonate (7.6mL, 50mmol) in anhydrous tetrahydrofuran (500mL) and place under an argon atmosphere. Cool to 5°C, add sodium hydride (1.2g, 50mmol) and stir briefly until homogeneous. Add  $\alpha,\alpha'$ -dibromo-o-xylene (13.2g, 50mmol) and stir an additional 15 minutes. Add additional sodium hydride (1.2g, 50mmol) and stir for 16 hours while warming to room temperature. Filter, evaporate the solvent in vacuo and purify by silica gel chromatography (1:1 methylene chloride/hexane) to give 2,2-dicarboethoxy-indan (9.64g, 74%).

Dissolve 2,2-dicarboethoxy-indan (5.67g, 21.7mmol) in ethanol (150mL). Add 1N lithium hydroxide (50mL) and stir overnight at room temperature. Reflux for 1 hour and concentrate the solution in vacuo. Partition between ethyl acetate and 6N hydrochloric acid. Separate the organic phase and wash with brine. Dry (MgSO<sub>4</sub>) and evaporate the solvent in vacuo to give an off-white solid. Distill (120-160°C @ 0.2-0.5 mmHg) to give 2-carboxy-indan (2.5g, 71%).

Dissolve 2-carboxy-indan (2.5g, 15.4mmol) in methanol and cool to 0°C. Saturate with hydrochloride gas then add 2,2-dimethoxypropane (2-3mL). Stir overnight then evaporate the solvent in vacuo. Purify by silica gel chromatography (2:1 methylene chloride/hexane) to give 2-carbomethoxy-indan as a water white oil. (2.04g, 75%).

Dissolve diisopropylamine (1.90mL, 7.8mmol) in anhydrous tetrahydrofuran (8mL), cool to -20°C and place under an argon atmosphere. Add, by dropwise addition, n-butyllithium (3.12mL of a 2.5N solution in hexanes, 7.8mmol) and stir for 20 minutes while cooling to -70°C. Add, by dropwise addition, a solution of 2-carbomethoxy-indan (1.349, 7.8mmol) in anhydrous tetrahydrofuran (8mL). Stir at -70°C for an additional 30 minutes then add trimethylsilyl chloride (freshly distilled from barium oxide, 1.0mL, 7.8mmol). Allow to warm to 10°C, evaporate the solvent in vacuo and dry the residue under vacuum for 30 minutes. Suspend the residue in methylene chloride (30mL), add zinc bromide (300mg, 1.3mmol) followed by t-butyl chloromethylsulfide (1.08g, 7.8mmol). Stir for 15 minutes at room temperature and add additional zinc bromide (500mg, 2.2mmol). Pour onto excess saturated sodium hydrogen carbonate and shake vigorously. Separate the organic phase and extract the aqueous phase with methylene chloride (30mL). Combine the organic phases, dry (MgSO<sub>4</sub>) and evaporate the solvent in vacuo to give crude product as a brown oil (2.12g, 98%). Purify by silica gel chromatography (30-70°C methylene chloride/hexane) and recrystallize (methanol) to give 2-carbomethoxy-2-(t-butyl)thiomethyl-indan as a crystalline solid (1.3g, 61%).

Dissolve 2-carbomethoxy-2-(t-butyl)thiomethyl-indan (557mg, 2.0mmol) in methanol (15mL) and add 1N lithium hydroxide (3.5mL). Warm briefly to effect solution then stir at room temperature under an argon atmosphere for 1 hour. Reflux for 6 hours, concentrate in vacuo to a volume of 3 mL and dilute to a volume of 15mL with water. Wash with methylene chloride and acidify and aqueous phase with excess 2N hydrochloric acid. After 5 minutes, collect the resulting white precipitate by filtration and dry to give 2-carboxy-2-(t-butyl)thiomethyl-indan (506mg, 96%); mp 158-163 °C.

Dissolve [4S-[4 $\alpha$ , 7 $\alpha$ (R), 12b $\beta$ ]]-7-(amino)-1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido[2,1-a][2]benzazepine-4-carboxylic acid, diphenylmethyl ester (150mg, 0.340mmol) and EDC (98mg, 0.51mmol) in tetrahydrofuran (5mL). Treat with a solution of 2-carboxy-2-(t-butyl)thiomethyl-indan (113mg, 0.426mmol). Stir at room temperature for 15 hours and evaporate the solvent in vacuo. Dissolve the residue in ethyl acetate (40mL) and wash with 1N hydrochloric acid (15mL) then saturated sodium hydrogen carbonate (15mL). Dry (Na<sub>2</sub>SO<sub>4</sub>) and evaporate the solvent in vacuo. Purify by silica gel chromatography (2:1 hexane/ethyl acetate) to give [4S-[4 $\alpha$ , 7 $\alpha$ (R), 12b $\beta$ ]]-7-[(2-(t-butylthio)methane-2-oxoindan)methylamino]-1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido[2,1-a][2]benzazepine-4-carboxylic acid, diphenylmethyl ester as a white solid (213mg, 91%).

Dissolve [4S-[4 $\alpha$ , 7 $\alpha$ (R), 12b $\beta$ ]]-7-[(2-(t-butylthio)methane-2-oxoindan)methylamino]-1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido[2,1-a][2]benzazepine-4-carboxylic acid, diphenylmethyl ester (210mg, 0.306mmol) in methylene chloride (4mL) and cool to 0 °C. Treat with trifluoroacetic acid (2mL), anisole (0.3mL, 2.7mmol), and mercuric acetate (122mg, 0.38mmol). Stir at room temperature overnight then bubble hydrogen sulfide gas through the solution for 10 minutes. Filter and dilute with methylene chloride. Wash organic phase with water (20mL), dry (Na<sub>2</sub>SO<sub>4</sub>) and evaporate the solvent in vacuo. Purify by silica gel chromatography (2.5:1 hexane/ethyl acetate to 1:1 hexane/ethyl acetate with 1% acetic acid) to give the title compound (144mg).

#### Example 9

[4S-[4 $\alpha$ , 7 $\alpha$ (R), 12b $\beta$ ]]-7-[(1-Thio-1-oxocyclopentane)methylamino]-1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido[2,1-a][2]benzazepine-4-carboxylic acid

Dissolve diisopropylamine (0.46mL, 3.3mmol) in tetrahydrofuran (10mL) and cool in an ice bath. Add, by dropwise addition, n-butyllithium (1.8mL of a 1.6M solution in hexanes, 2.9mmol). Stir for 15 minutes, cool to -78 °C and add a solution of methyl cyclopentanecarboxylate (322mg, 2.51mmol) in tetrahydrofuran (5mL). Stir for 1 hour then treat with hexamethylphosphoramide (0.66mL, 3.8mmol). Stir for 15 minutes then add 4-methoxybenzyl disulfide (1.0g, 3.3mmol) in tetrahydrofuran (13mL). Stir for 3 hours at -78 °C, quench with saturated aqueous ammonium chloride (5mL). Partition between water (2X50mL) and ethyl acetate (100mL). Dry (Na<sub>2</sub>SO<sub>4</sub>) and pass through a plug of silica gel (methylene chloride) to give 1-(carbomethoxy)-1-(4-methoxybenzylthio)cyclopentane as a pale yellow oil.

Dissolve 1-(carbomethoxy)-1-(4-methoxybenzylthio)cyclopentane (3.3mmol) in 95% ethanol (18mL), water (9mL), and tetrahydrofuran (10mL). Treat with potassium hydroxide (0.91g, 16mmol). Stir at room temperature for 2 hours and evaporate the solvent in vacuo. Partition between water (50mL) and ethyl ether (30mL). Acidify the aqueous phase with cold concentrated hydrochloric acid and extract with methylene chloride (50mL). Dry (MgSO<sub>4</sub>) and evaporate the solvent in vacuo to give 1-(carboxy)-1-(4-methoxybenzylthio)cyclopentane as a pale yellow oil (483mg, 72%).

Dissolve [4S-[4 $\alpha$ , 7 $\alpha$ (R), 12b $\beta$ ]]-7-(amino)-1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido[2,1-a][2]benzazepine-4-carboxylic acid, diphenylmethyl ester (200mg, 0.454mmol), EDC (130mg, 0.68mmol) and 1-(carboxy)-1-(4-methoxybenzylthio)cyclopentane (128mg, 0.480mmol) in tetrahydrofuran (5mL). Stir at room temperature for 16 hours and evaporate the solvent in vacuo. Dissolve the residue in ethyl acetate (30mL) and wash with 5% sulfuric acid (15mL) then saturated sodium hydrogen carbonate (15mL). Dry (Na<sub>2</sub>SO<sub>4</sub>) and evaporate the solvent in vacuo. Purify by silica gel chromatography (2:1 hexane/ethyl acetate to 3:2 hexane/ethyl acetate) to give [4S-[4 $\alpha$ , 7 $\alpha$ (R), 12b $\beta$ ]]-7-[(1-(4-methoxybenzylthio)-1-oxo-cyclopentane)methylamino]-1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido[2,1-a][2]benzazepine-4-carboxylic acid, diphenylmethyl ester as a white solid (195mg, 62.3%).

Dissolve [4S-[4 $\alpha$ , 7 $\alpha$ (R), 12b $\beta$ ]]-7-[(1-(4-methoxybenzylthio)-1-oxo-cyclopentane)methylamino]-1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido[2,1-a][2]benzazepine-4-carboxylic acid, diphenylmethyl ester (195mg, 0.283mmol) in methylene chloride (4mL) and cool to 0 °C. Treat with trifluoroacetic acid (2mL), anisole (0.3mL, 2.7mmol), and mercuric acetate (113mg, 0.354mmol). Stir for 3 hours then bubble hydrogen sulfide gas through the solution for 15 minutes. Filter and dilute with methylene chloride. Wash organic phase with water (20mL), dry (Na<sub>2</sub>SO<sub>4</sub>) and evaporate the solvent in vacuo. Purify by silica gel chromatography (2:1 hexane/ethyl acetate to 1:1 hexane/ethyl acetate with 1% acetic acid) to give the title compound

(110mg, 96.5%).

#### Example 10

5 [4S-{4 $\alpha$ , 7 $\alpha$ (R'), 12b $\beta$ }]-7-[(2-Thio-1-oxo-2-methyl)-2-propylamino]-1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido-  
[2,1-a][2]benzazepine-4-carboxylic acid

Dissolve 4-methoxybenzylthiol (14mL, 0.10mmol) in anhydrous dimethylformamide (150mL), degas and purge with nitrogen. Add diisopropylethylamine (20ml, 0.115mol), then treat with ethyl bromoacetate  
 10 (11.1mL, 0.10mol). Place the reaction flask in a cooling bath containing cold water and stir to room temperature for 64 hours. Partition between water (300mL) and ether (250mL). Separate the aqueous phase and extract with ether (250mL). Combine the organic phases, wash with water (2X150mL) and extract the combined aqueous phases with ether (125mL). Dry (Na<sub>2</sub>SO<sub>4</sub>) and evaporate the solvent *in vacuo*. Purify by chromatography (6:1 hexane/ethyl acetate) to give 1-(carboethoxy)-1-(4-methoxybenzylthio)methane as a  
 15 yellow oil (22g, 92%).

Dissolve lithium hexamethyldisilazane (4.2mL of a 1.0M solution in tetrahydrofuran, 4.2mmol) in tetrahydrofuran (9mL) and cool to -78°C. Treat with a solution of 1-(carboethoxy)-1-(4-methoxybenzylthio)-  
 20 methane (1.0g, 4.2mmol) in tetrahydrofuran (4mL). Stir for 45 minutes, add methyl iodide (0.26mL, 4.2mmol) and stir for 3 hours. Add additional lithium hexamethyldisilazane (4.2mL of a 1.0M solution in tetrahydrofuran, 4.2mmol), stir for 30 minutes and add methyl iodide (0.3mL, 4.8mmol). Stir at room temperature overnight. Partition between saturated ammonium chloride (30mL) and ethyl acetate (50mL). Separate the organic phase, dry (Na<sub>2</sub>SO<sub>4</sub>) and evaporate the solvent *in vacuo*. Purify by chromatography (5:1 hexane/ethyl acetate) to give 2-(carboethoxy)-2-(4-methoxybenzylthio)propane as a pale yellow oil  
 (1.05g, 93.8%).

25 Dissolve 2-(carboethoxy)-2-(4-methoxybenzylthio)propane (1.05g, 3.91mmol) in a mixture of 95% ethanol (18mL), water (9mL) and tetrahydrofuran (10mL). Add potassium hydroxide (1.49, 25mmol) and stir at room temperature for 2 hours then at 55°C for 1 hour. Evaporate the solvent *in vacuo* and partition between water (50mL) and ether (50mL). Acidify to pH 1 with concentrated hydrochloric acid and extract into methylene chloride (50mL). Dry (Na<sub>2</sub>SO<sub>4</sub>) and evaporate the solvent *in vacuo* to give 2-(carboxy)-2-(4-  
 30 methoxybenzylthio)propane as a tan solid (763mg).

Dissolve [4S-{4 $\alpha$ , 7 $\alpha$ (R'), 12b $\beta$ }]-7-(amino)-1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido[2,1-a][2]benzazepine-4-  
 carboxylic acid, diphenylmethyl ester (175mg, 0.40mmol, EEDQ (101mg, 0.41mmol) and 2-(carboxy)-2-(4-  
 35 methoxybenzylthio)propane (96mg, 0.40mmol in methylene chloride (5mL). Stir at room temperature for 18 hours and evaporate the solvent *in vacuo*. Dissolve the residue in ethyl acetate (30mL) and wash with 5% sulfuric acid (15ml) then saturated sodium hydrogen carbonate (15mL). Dry (Na<sub>2</sub>SO<sub>4</sub>) and evaporate the solvent *in vacuo*. Purify by silica gel chromatography (3:2 hexane/ethyl acetate to 1:1 hexane/ethyl acetate) to give  
 [4S-{4 $\alpha$ , 7 $\alpha$ (R'), 12b $\beta$ }]-7-[(2-(4-methoxybenzylthio)-1-oxo-2-methyl)-2-propylamino]-  
 1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido[2,1-a][2]benzazepine-4-carboxylic acid, diphenylmethyl ester as a  
 white foam (260mg).

40 Dissolve [4S-{4 $\alpha$ , 7 $\alpha$ (R'), 12b $\beta$ }]-7-[(2-(4-methoxybenzylthio)-1-oxo-2-methyl)-2-propylamino]-  
 1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido[2,1-a][2]benzazepine-4-carboxylic acid, diphenylmethyl ester  
 (260mg, 0.40mmol) in methylene chloride (5mL) and cool to 0°C. Treat with anisole (0.43mL, 4.0mmol) and mercuric acetate (159mg, 0.50mmol) then with trifluoroacetic acid (2.5mL). Stir for 3 hours then bubble  
 45 hydrogen sulfide gas through the solution for 10 minutes. Filter and dilute with methylene chloride. Wash organic phase with water (20mL), dry (Na<sub>2</sub>SO<sub>4</sub>) and evaporate the solvent *in vacuo*. Purify by silica gel chromatography (1:1 hexane/ethyl acetate with 1% acetic acid) to give the title compound as a white solid  
 (114mg).

#### Example 11

50 [4S-{4 $\alpha$ , 7 $\alpha$ (R'), 12b $\beta$ }]1-7-[(1-Thio-1-oxocyclopropane)methylamino]-1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido-  
[2,1-a][2]benzazepine-4-carboxylic acid

Cool lithium hexamethyldisilazane (6mL of a 1.0M solution in tetrahydrofuran, 6.0mmol) to -78°C. Add, by  
 55 cannula transfer, a solution of 1-(carboethoxy)-1-(4-methoxybenzylthio)methane (1.36g, 5.66mmol) in tetrahydrofuran (10mL). Stir for 45 minutes, add 1,3-dimethyl-3,4,5,6-tetrahydro-2(1H)-pyrimidinone (DMPU) (0.73mL, 6.0mmol) followed by 1-bromo-2-chloroethane (0.50mL, 6.0mmol) and stir to room temperature overnight. Cool to -78°C, add lithium hexamethyldisilazane (6.0mL, 6.0mmol) and stir at room temperature

for 24 hours. Quench with saturated ammonium chloride (5mL) and partition between ethyl acetate (30mL) and water (30mL). Separate the organic phase, dry ( $\text{Na}_2\text{SO}_4$ ) and evaporate the solvent *in vacuo*. Purify by chromatography (5:1 hexane/ethyl acetate) to 2.5:1 hexane/ethyl acetate) to give 1-(carboethoxy)-1-(4-methoxybenzylthio)cyclopropane (317mg, 21%).

- 5 Dissolve 1-(carboethoxy)-1-(4-methoxybenzylthio)cyclopropane (0.310g, 1.16mmol) in 95% ethanol (6mL), water (3mL), and tetrahydrofuran (6mL). Treat with lithium hydroxide monohydrate (420mg, 10mmol). Stir at a gentle reflux for 6 hours and evaporate the solvent *in vacuo*. Add water (50mL) and wash with ether (50mL). Acidify the aqueous phase and extract into methylene chloride (50mL). Dry ( $\text{Na}_2\text{SO}_4$ ) and evaporate the solvent *in vacuo* to give 1-(carboxy)-1-(4-methoxybenzylthio)cyclopropane as a pale yellow solid (266mg).

- 10 Dissolve [4S-[4 $\alpha$ , 7 $\alpha$ (R\*), 12b $\beta$ ]]-7-(amino)-1,2,3,5,6,7,8,12b-octahydro-6-oxopyrido[2,1-a][2]benzazepine-4-carboxylic acid, diphenylmethyl ester (220mg, 0.50mmol), EDC (144mg, 0.75mmol) and 1-(carboxy)-1-(4-methoxybenzylthio)cyclopropane (120mg, 0.50mmol) in tetrahydrofuran (5mL). Stir at room temperature for 24 hours and evaporate the solvent *in vacuo*. Dissolve the residue in ethyl acetate (30mL) and wash with 5% sulfuric acid (20mL), saturated sodium hydrogen carbonate (15mL) and brine (15mL). Dry ( $\text{Na}_2\text{SO}_4$ ) and evaporate the solvent *in vacuo*. Purify by silica gel chromatography (3:2 hexane/ethyl acetate to 2:3 hexane/ethyl acetate) to give [4S-[4 $\alpha$ , 7 $\alpha$ (R\*), 12b $\beta$ ]]-7-[(1-(4-methoxybenzylthio)-1-oxo-cyclopropane)-methylamino]-1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido[2,1-a][2]benzazepine-4-carboxylic acid, diphenylmethyl ester (128mg, 39%).

- 20 Dissolve [4S-[4 $\alpha$ , 7 $\alpha$ (R\*), 12b $\beta$ ]]-7-[(1-(4-methoxybenzylthio)-1-oxo-cyclopropane)methylamino]-1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido[2,1-a][2]benzazepine-4-carboxylic acid, diphenylmethyl ester (128mg, 0.193mmol) in methylene chloride (3mL) and add anisole (0.21mL, 1.94mmol) and mercuric acetate (78mg, 0.245mmol). Cool to 0°C and treat with trifluoroacetic acid (1.5mL). Stir for 3.5 hours then bubble hydrogen sulfide gas through the solution for 15 minutes. Filter, evaporate the solvent *in vacuo* and dilute with methylene chloride and carbon tetrachloride. Evaporate the solvent *in vacuo* and purify by silica gel chromatography (1:1 hexane/ethyl acetate with 1% acetic acid) to give the title compound as a white solid (70mg).

#### Example 12

- 30 [4S-[4 $\alpha$ , 7 $\alpha$ (R\*), 12b $\beta$ ]]-7-[(1-Thio-1-oxocyclohexane)methylamino]-1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido[2,1-a][2]benzazepine-4-carboxylic acid

- Cool lithium hexamethyldisilazane (8.4mL of a 1.0M solution in tetrahydrofuran, 8.4mmol) to -78°C. Add, by cannula transfer, a solution of 1-(carboethoxy)-1-(4-methoxybenzylthio)methane (2.0g, 8.3mmol) in tetrahydrofuran (8mL). Stir for 45 minutes then add a solution of 1-iodo-5-chloropentane (1.94g, 8.32mmol) in tetrahydrofuran (8mL). Stir for 1 hour and add 1,3-dimethyl-3,4,5,6-tetrahydro-2(1H)-pyrimidinone (DMPU) (1.21mL, 10.0mmol). Stir at 0°C for 4 hours and at room temperature for 6 hours. Cool to -78°C, add lithium hexamethyldisilazane (8.4mL, 8.4mmol) and stir at room temperature overnight. Quench with saturated ammonium chloride (5mL) and partition between ethyl acetate (50mL) and water (50mL). Separate the organic phase, dry ( $\text{Na}_2\text{SO}_4$ ) and evaporate the solvent *in vacuo*. Purify by chromatography (6:1 hexane/ethyl acetate) to give 1-(carboethoxy)-1-(4-methoxybenzylthio)cyclohexane as a colorless oil (1.45g, 56%).

- Dissolve 1-(carboethoxy)-1-(4-methoxybenzylthio)cyclohexane (1.45g, 4.7mmol) in water (2.5mL) and tetrahydrofuran (10mL). Treat with 2.5N sodium hydroxide (4mL, 10mmol).

- Heat at 60°C for 2 hours, add 95% ethanol (10mL) and lithium hydroxide monohydrate (0.42g, 10mmol). Heat to a gentle reflux for 20 hours, cool and evaporate the solvent *in vacuo*. Partition between water (50mL) and ether (50mL). Separate the aqueous phase and acidify to pH 1. Extract with methylene chloride (50mL), dry ( $\text{Na}_2\text{SO}_4$ ) and evaporate the solvent *in vacuo* to give 1-(carboxy)-1-(4-methoxybenzylthio)-cyclohexane as a white solid (1.21g, 92%).

- 50 Dissolve [4S-[4 $\alpha$ , 7 $\alpha$ (R\*), 12b $\beta$ ]]-7-(amino)-1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido[2,1-a][2]benzazepine-4-carboxylic acid, diphenylmethyl ester (220mg, 0.50mmol), EDC (144mg, 0.75mmol) and 1-(carboxy)-1-(4-methoxybenzylthio)cyclohexane (140mg, 0.50mmol) in tetrahydrofuran (6mL). Stir at room temperature for 24 hours, add additional EDC (95mg, 0.5mmol) and 1-(carboxy)-1-(4-methoxybenzylthio)cyclohexane (70mg, 0.25mmol). Stir at room temperature overnight and evaporate the solvent *in vacuo*. Dissolve the residue in ethyl acetate (30mL) and wash with 5% sulfuric acid (15mL), saturated sodium hydrogen carbonate (15mL) and brine (15mL). Dry ( $\text{Na}_2\text{SO}_4$ ) and evaporate the solvent *in vacuo*. Purify by silica gel chromatography (2:1 hexane/ethyl acetate) to give [4S-[4 $\alpha$ , 7 $\alpha$ (R\*), 12b $\beta$ ]]-7-[(1-(4-methoxybenzylthio)-1-oxo-cyclohexane)-

methylamino]-1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido[2,1-a][2]benzazepine-4-carboxylic acid, diphenylmethyl ester (220mg, 63%).

- Dissolve [4S-[4 $\alpha$ , 7 $\alpha$ (R\*), 12b $\beta$ ]]-7-[(1-(4-methoxybenzylthio)-1-oxo-cyclohexane)methylamino]-1,2,3,5,6,7,8,12b-octahydro-6-oxopyrido[2,1-a][2]benzazepine-4-carboxylic acid, diphenylmethyl ester (220mg, 0.313mmol) in methylene chloride (5mL) and add anisole (0.34mL, 3.13mmol) and mercuric acetate (125mg, 0.39mmol). Cool to 0 °C and treat with trifluoroacetic acid (2.5mL). Stir for 3 hours then bubble hydrogen sulfide gas through the solution for 10 minutes. Filter, evaporate the solvent *in vacuo*. Dilute with methylene chloride and carbon tetrachloride and evaporate the solvent *in vacuo* again. Purify by silica gel chromatography (1:1 hexane/ethyl acetate with 1% acetic acid) to give the title compound as a white solid (100mg, 77%).

### Example 13

- [4S-[4 $\alpha$ , 7 $\alpha$ (R\*), 12b $\beta$ ]]-7-[(5-Thio-5-oxo-4,5-dihydrocyclopentimidazole)methylamino]-3,4,6,7,8,12b-hexahydro-6-oxo-1H-[1,4]-oxazino[3,4-a][2]benzazepine-4-carboxylic acid

Scheme D, step a: N-[2-(1,3-dihydro-1,3-dioxo-2H-isoindol-2-yl)-1-oxo-3-phenylpropyl]-L-serine, methyl ester

- Slurry N-phthaloyl-(S)-phenylalanine (90g, 0.3mol) in methylene chloride (450mL) and add, by dropwise addition, oxalyl chloride (54mL, 0.62mol). Place under a dry atmosphere (CaSO<sub>4</sub> tube) and treat with dimethylformamide (10 $\mu$ L). Stir for 5 hours, filter and concentrate *in vacuo* to give N-phthaloyl-(S)-phenylalanine, acid chloride an off white amorphous solid. Dissolve serine methyl ester hydrochloride (56g, 0.36mol) in tetrahydrofuran (300mL) then cool to 0 °C and add 4-methylmorpholine (88mL, 0.8mol). Add, by dropwise addition, a solution of the N-phthaloyl-(S)-phenylalanine, acid chloride in tetrahydrofuran (200mL). Allow to warm to room temperature and stir for 3 hours. Filter and concentrate the filtrate *in vacuo*. Dissolve the residue in ethyl acetate and separate the organic phase. Wash with water then saturated sodium chloride and dry (MgSO<sub>4</sub>). Evaporate the solvent *in vacuo* to give an oil. Purify by silica gel chromatography (gradient 50% ethyl acetate/hexane to ethyl acetate) to give the title compound (80.8g, 67%) mp 129-132 °C.

Scheme D, step b: N-[2-(1,3-Dihydro-1,3-dioxo-2H-isoindol-2-yl)-1-oxo-3-phenylpropyl]-O-2-propenyl-L-serine, methyl ester

- Dissolve N-[2-(1,3-dihydro-1,3-dioxo-2H-isoindol-2-yl)-1-oxo-3-phenylpropyl]-L-serine, methyl ester (25g, 63mmol) in methylene chloride/cyclohexane (1:1, 600mL). Add allyl trichloroacetimidate (26g, 128mmol) and trifluoromethanesulfonic acid (5mL, 56.6mmol). Stir at room temperature under a nitrogen atmosphere for 5 hours and dilute with methylene chloride. Wash with saturated aqueous sodium hydrogen carbonate, water, dry (MgSO<sub>4</sub>) and evaporate the solvent *in vacuo*. Purify by silica gel chromatography (gradient 20% ethyl acetate/hexane to 35% ethyl acetate/hexane) to give the title compound; mp 95-97 °C.

Scheme D, step c: [S-(R\*, R\*)]-N-[2-(1,3-Dihydro-1,3-dioxo-2H-isoindol-2-yl)-1-oxo-3-phenylpropyl]-3,4-dihydro-2H-1,4-oxazine-3-carboxylic acid, methyl ester

- Dissolve N-[2-(1,3-dihydro-1,3-dioxo-2H-isoindol-2-yl)-1-oxo-3-phenylpropyl]-O-2-propenyl-L-serine, methyl ester (13g, 29.8mmol) in methylene chloride/methanol (10:1, 220mL). Cool to -78 °C and sparge with a mixture of ozone/oxygen for approximately 10 minutes until a blue color persists. Sparge with nitrogen for 10 minutes at -78 °C to remove excess ozone. Treat with methylsulfide (60mL, 0.82mol) and allow to warm to room temperature. Stir at room temperature for 2.5 hours, evaporate the solvent *in vacuo* and dissolve the residue in ethyl acetate (200mL). Wash with water, saturated sodium chloride, dry (MgSO<sub>4</sub>) and evaporate the solvent *in vacuo* to give the intermediate N-[2-(1,3-dihydro-1,3-dioxo-2H-isoindol-2-yl)-1-oxo-3-phenylpropyl]-O-2-oxoethyl-L-serine, methyl ester as a foam (13.6g). Dissolve N-[2-(1,3-dihydro-1,3-dioxo-2H-isoindol-2-yl)-1-oxo-3-phenylpropyl]-O-2-oxoethyl-L-serine, methyl ester (13.6g) in methylene chloride/trifluoroacetic acid (10:1/330mL). Stir at room temperature for 2.5 hours and evaporate the solvent *in vacuo*. Purify by silica gel chromatography (35% ethyl acetate/hexane) and recrystallize (ethyl acetate/hexane) to give the title compound (8.52g, 68%); mp 70-72 °C.



Scheme D, step d: [4S-[4 $\alpha$ , 7 $\alpha$ (R'), 12b $\beta$ ]]-7-[(1,3-Dihydro-1,3-dioxo-2H-isoindol-2-yl)]-3,4,6,7,8,12b-hexahydro-6-oxo-1H-[1,4]-oxazino[3,4-a][2]benzazepine-4-carboxylic acid, diphenylmethyl ester

Dissolve [S-(R', R')]-N-[2-(1,3-dihydro-1,3-dioxo-2H-isoindol-2-yl)-1-oxo-3-phenylpropyl]-3,4-dihydro-2H-1,4-oxazine-3-carboxylic acid, methyl ester (2.5g, 5.9mmol) in methylene chloride (5mL) and add, by dropwise addition, to a previously prepared solution of trifluoromethanesulfonic acid (4.0mL, 45mmol) and trifluoroacetic anhydride (1.0mL, 7.1mmol). Place under a nitrogen atmosphere and stir at room temperature for 123 hours. Pour into a separatory funnel containing ice (200g) and ethyl acetate (200mL). Separate the organic phase, wash with water (3X200mL) and saturated aqueous sodium chloride (100mL). Extract the organic phase with 10% wt. potassium hydrogen carbonate (4X40mL) and water (40mL). Layer the combined basic aqueous phases with ethyl acetate (100mL) and cool in an ice bath. Add, by dropwise addition, 6N hydrochloric acid to adjust the pH to 1 while maintaining the temperature at 5-10 °C. Separate the organic phase and extract the aqueous phase with ethyl acetate (3X200mL), wash with saturated sodium chloride and dry (MgSO<sub>4</sub>). Evaporate the solvent in vacuo and dry the residue under high vacuum at 56 °C for 24 hours to give the intermediate [4S-[4 $\alpha$ , 7 $\alpha$ (R'), 12b $\beta$ ]]-7-[(1,3-dihydro-1,3-dioxo-2H-isoindol-2-yl)]-3,4,6,7,8,12b-hexahydro-6-oxo-1H-[1,4]-oxazino[3,4-a][2]benzazepine-4-carboxylic acid (1.75g, 73%). Dissolve [4S-[4 $\alpha$ , 7 $\alpha$ (R'), 12b $\beta$ ]]-7-[(1,3-dihydro-1,3-dioxo-2H-isoindol-2-yl)]-3,4,6,7,8,12b-hexahydro-6-oxo-1H-[1,4]-oxazino[3,4-a][2]benzazepine-4-carboxylic acid (500mg, 1.23mmol) in methylene chloride (12mL) and treat with diphenyldiazomethane (360mg, 1.86mmol). Stir for 5.5 hours and evaporate the solvent in vacuo. Purify by silica gel chromatography (gradient 20% ethyl acetate/hexane to 35% ethyl acetate/hexane) to give the title compound (563mg, 80%); mp 178-181 °C (isopropanol).

Scheme D, step e: [4S-[4 $\alpha$ , 7 $\alpha$ (R'), 12b $\beta$ ]]-7-(Amino)-3,4,6,7,8,12b-hexahydro-6-oxo-1H-[1,4]-oxazino[3,4-a][2]benzazepine-4-carboxylic acid, diphenylmethyl ester

Dissolve [4S-[4 $\alpha$ , 7 $\alpha$ (R'), 12b $\beta$ ]]-7-[(1,3-dihydro-1,3-dioxo-2H-isoindol-2-yl)]-3,4,6,7,8,12b-hexahydro-6-oxo-1H-[1,4]-oxazino[3,4-a][2]benzazepine-4-carboxylic acid, diphenylmethyl ester (296mg, 0.517mmol) in methanol (5mL) and treat with hydrazine monohydrate (1.1mL of a 1M solution in methanol, 1.1mmol). Stir at room temperature for 44 hours, evaporate the solvent in vacuo and slurry the residue in methylene chloride (10mL). Filter and evaporate the solvent in vacuo to give the title compound (218mg, 95%).

Scheme A: [4S-[4 $\alpha$ , 7 $\alpha$ (R'), 12b $\beta$ ]]-7-[(5-Thio-5-oxo-4,5-dihydro-cyclopentimidazole)methylamino]-3,4,6,7,8,12b-hexahydro-6-oxo-1H-[1,4]-oxazino[3,4-a][2]benzazepine-4-carboxylic acid

Dissolve 4,5-imidazoledicarboxylic acid (31.2g, 0.2mol) in ethanol (500mL) and treat with concentrated sulfuric acid (0.5mL). Heat to 60 °C for 16 hours, cool and reduce the solvent by 50% in vacuo. Dilute with ethyl ether (500mL), wash with saturated sodium hydrogen carbonate, then brine. Dry (MgSO<sub>4</sub>) and evaporate the solvent in vacuo to give 4,5-(dicarboethoxy)imidazole. Dissolve 4,5-(dicarboethoxy)imidazole (1.06g, 5mmol) and triethylamine (1.5mL, 7.5mmol) in 50/50 dioxane water (25mL). Add [2-(tert-butyloxycarbonyloxymino)-2-phenylacetone nitrile] (1.36g, 5.5mmol) and stir at room temperature for 2 hours. Add water (7.5mL) and ethyl acetate (10mL), separate the aqueous phase and wash with ethyl acetate (10mL). Combine the organic phases, dry (MgSO<sub>4</sub>) and evaporate the solvent in vacuo. Purify the residue by silica gel chromatography to give N-(carbo-t-butyloxy)-4,5-(dicarboethoxy)imidazole. Dissolve N-(carbo-t-butyloxy)-4,5-(dicarboethoxy)imidazole (3.12g, 10mmol) in anhydrous tetrahydrofuran (30mL) and cool to -20 °C. Treat with lithium borohydride (7mL of a 2N solution) and stir under a nitrogen atmosphere for several days. Carefully add water and partition between ethyl acetate and 5% hydrochloric acid. Separate the organic phase, wash with brine and dry (MgSO<sub>4</sub>). Evaporate the solvent in vacuo and purify by silica gel chromatography to give N-(carbo-t-butyloxy)-4,5-(dihydroxymethyl)imidazole. Dissolve N-bromosuccinimide (1.78g, 0.01mol) in tetrahydrofuran (60mL) and add a solution of triphenylphosphine (2.62g, 0.01mol) in tetrahydrofuran. Add a solution of N-(carbo-t-butyloxy)-4,5-(dihydroxymethyl)imidazole (1.14g, 5mmol) in tetrahydrofuran (25mL) and stir until most of the solid goes into solution. Evaporate the solvent in vacuo and partition the residue between water and ethyl ether. Separate the organic phase and wash with water. Dry (Na<sub>2</sub>SO<sub>4</sub>) and evaporate the solvent in vacuo. Purify by silica gel chromatography to give N-(carbo-t-butyloxy)-4,5-(dibromomethyl)imidazole. Dissolve diethylmalonate (15.2mL, 0.100mol) in tetrahydrofuran (800mL). Cool in an ice bath and treat with sodium hydride (3.0g, 0.10mol, 80% in mineral oil). Stir until solution attained and add N-(carbo-t-butyloxy)-4,5-(dibromomethyl)imidazole (35.4g, 0.100mol). Stir for 30 minutes then add additional sodium hydride

(3.0g, 0.10mol). Stir at room temperature for 20 hours, filter through filter aid and evaporate the solvent in vacuo. Purify by silica gel chromatography to give 5,5-(dicarboethoxy)-1-(carbo-t-butyloxy)-4,5-dihydro-cyclopentimidazole.

Dissolve 5,5-(dicarboethoxy)-1-(carbo-t-butyloxy)-4,5-dihydro-cyclopentimidazole (21.3g, 60.6mmol) in dimethylsulfoxide (140mL). Add water (14mL) and lithium chloride (7.0g, 0.16mol). Heat at reflux for 4 hours, cool and partition between water (150mL) and methylene chloride (2X150mL). Wash the organic phase with water (150mL), dry (MgSO<sub>4</sub>) and pass through a silica gel plug to give 5-(carboethoxy)-1-(carbo-t-butyloxy)-4,5-dihydro-cyclopentimidazole.

Dissolve 5-(carboethoxy)-1-(carbo-t-butyloxy)-4,5-dihydro-cyclopentimidazole (9.5g, 34.1mmol) in ethanol (95%, 150mL) and water (75mL). Add potassium hydroxide (9.5g, 0.17mol) and stir at room temperature for 1 hour. Partition between water (150mL) and ethyl ether (2X150mL). Acidify the aqueous phase with hydrochloric acid to pH 1. Extract with methylene chloride (2X150mL), dry (Na<sub>2</sub>SO<sub>4</sub>) and evaporate the solvent in vacuo to give 5-(carboxy)-1-(carbo-t-butyloxy)-4,5-dihydro-cyclopentimidazole.

Dissolve 5-(carboxy)-1-(carbo-t-butyloxy)-4,5-dihydro-cyclopentimidazole (5.9g, 23.5mmol) in methanol (60mL) and treat with dimethoxypropane (5.8mL, 47mmol) and sulfuric acid (0.8mL). Stir at room temperature for 1 day. Evaporate the solvent in vacuo, dilute with methylene chloride (75mL) and wash with saturated sodium hydrogen carbonate (35mL). Extract the aqueous phase with methylene chloride (30mL), wash combined organics with brine (30mL) and dry (Na<sub>2</sub>SO<sub>4</sub>). Evaporate the solvent in vacuo and pass through a plug of silica gel to give 5-(carbomethoxy)-1-(carbo-t-butyloxy)-4,5-dihydro-cyclopentimidazole.

Mix 4-methoxybenzylthiol (3.0g, 19mmol) in sodium hydroxide (20mL of a 2.5N aqueous solution) and methanol (10mL). Add saturated aqueous copper sulfate solution (1.5mL) and stir at room temperature for 2 hours, blowing air over the top of the mixture. Filter, wash solid with water and dry to give 4-methoxybenzyl disulfide as a pale yellow powder (2.71g, 91%).

Cool lithium hexamethyldisilazane (4.2mL, 4.2mol, 1.0M in tetrahydrofuran) to -78°C and treat with a solution of 5-(carbomethoxy)-1-(carbo-t-butyloxy)-4,5-dihydro-cyclopentimidazole (944mg, 3.55mmol) in tetrahydrofuran (5mL). Stir for 1 hour then add hexamethylphosphoramide (0.93mL, 5.3mmol) and stir for 5 minutes. Add 4-methoxybenzyl disulfide (1.6g, 5.2mmol) in tetrahydrofuran (10mL). Stir for 5 hours at -78°C and quench with a solution of ammonium chloride. Partition between ethyl acetate (75mL) and brine (30mL). Dry (Na<sub>2</sub>SO<sub>4</sub>), evaporate the solvent in vacuo and purify by silica gel chromatography to give 5-(carbomethoxy)-1-(carbo-t-butyloxy)-5-(4-methoxybenzylthio)-4,5-dihydro-cyclopentimidazole.

Dissolve 5-(carbomethoxy)-1-(carbo-t-butyloxy)-5-(4-methoxybenzylthio)-4,5-dihydro-cyclopentimidazole (1.43g, 3.55mmol) in 95% ethanol (25mL), water (12mL) and tetrahydrofuran (15mL). Treat with potassium hydroxide (1.3g, 23mmol) and stir at room temperature for 1 hour. Filter and evaporate the solvent in vacuo. Partition between water (125mL) and ether (75mL). Separate the aqueous phase and acidify with cold concentrated hydrochloric acid. Extract with methylene chloride (75mL), dry (Na<sub>2</sub>SO<sub>4</sub>) and evaporate the solvent in vacuo. Purify by silica gel chromatography to give 5-(carboxy)-1-(carbo-t-butyloxy)-5-(4-methoxybenzylthio)-4,5-dihydro-cyclopentimidazole.

Dissolve [4S-[4 $\alpha$ , 7 $\alpha$ (R'), 12b $\beta$ ]]-7-(amino)-3,4,6,7,8,12b-hexahydro-6-oxo-1H-[1,4]-oxazino[3,4-a][2]-benzazepine-4-carboxylic acid, diphenylmethyl ester (201mg, 0.454mmol) and EDC (130mg, 0.68mmol) in tetrahydrofuran (5mL). Treat with 5-(carboxy)-1-(carbo-t-butyloxy)-5-(4-methoxybenzylthio)-4,5-dihydro-cyclopentimidazole (177mg, 0.454mmol). Stir at room temperature for 20 hours and evaporate the solvent in vacuo. Dissolve the residue in ethyl acetate (50mL) and wash with 5% sulfuric acid (15mL) then saturated sodium hydrogen carbonate (15mL). Dry (Na<sub>2</sub>SO<sub>4</sub>) and evaporate the solvent in vacuo. Purify by silica gel chromatography to give [4S-[4 $\alpha$ , 7 $\alpha$ (R'), 12b $\beta$ ]]-7-[(5-(4-methoxybenzylthio)-5-oxo-1-(carbo-t-butyloxy)-4,5-dihydro-cyclopentimidazole)methylamino]-3,4,6,7,8,12b-hexahydro-6-oxo-1H-[1,4]-oxazino[3,4-a][2]-benzazepine-4-carboxylic acid, diphenylmethyl ester.

Dissolve [4S-[4 $\alpha$ , 7 $\alpha$ (R'), 12b $\beta$ ]]-7-[(5-(4-methoxybenzylthio)-5-oxo-1-(carbo-t-butyloxy)-4,5-dihydro-cyclopentimidazole)methylamino]-3,4,6,7,8,12b-hexahydro-6-oxo-1H-[1,4]-oxazino[3,4-a][2]-benzazepine-4-carboxylic acid, diphenylmethyl ester (133mg, 0.163mmol) in methylene chloride (3mL) and cool to 0°C. Treat with trifluoroacetic acid (1.5mL), anisole (0.19mL, 1.7mmol), and mercuric acetate (65mg, 0.2mmol). Stir at 0°C for 3 hours then bubble hydrogen sulfide gas through the solution for 10 minutes. Filter and wash with methylene chloride. Wash organic phase with water (20mL), dry (Na<sub>2</sub>SO<sub>4</sub>) and evaporate the solvent in vacuo. Purify by silica gel chromatography to give the title compound.

## Example 14

Preparation of [4S-[4 $\alpha$ , 7 $\alpha$ (R'), 12b $\beta$ ]]-7-[(5-Thio-5-oxo-4,5,6-trihydro-cyclopenta[c]furan)methylamino]-3,4,6,7,8,12b-hexahydro-6-oxo-1H-[1,4]-thiazino[3,4-a][2]benzazepine-4-carboxylic acid

- 5 Dissolve 3,4-(dicarboethoxy)furan (21.2g, 10mmol) in anhydrous tetrahydrofuran (30mL) and cool to -20 °C. Treat with lithium borohydride (7mL of a 2N solution) and stir under a nitrogen atmosphere for several days. Carefully add water and partition between ethyl acetate and 5% hydrochloric acid. Separate the organic phase, wash with brine and dry (MgSO<sub>4</sub>). Evaporate the solvent in vacuo and purify by silica gel chromatography to give 3,4-(dihydroxymethyl)furan.
- 10 Dissolve 3,4-(dihydroxymethyl)furan (11.9g, 0.093mol) in 2,4,6-collidine (12.4g) and add a solution of lithium bromide (7.9g, 91mmol) in dimethylformamide (80mL). Cool to -50 °C and place under a nitrogen atmosphere. Add methanesulfonyl chloride (11.8g) at a rate which keeps the temperature under 0 °C. Stir at 0 °C for 2 hours and pour onto ice water. Extract with ethyl ether, dry (MgSO<sub>4</sub>), evaporate the solvent in vacuo and purify by silica gel chromatography to give 3,4-(dibromomethyl)furan.
- 15 Dissolve diethylmalonate (15.2mL, 0.100mol) in tetrahydrofuran (800mL). Cool in an ice bath and treat with sodium hydride (3.0g, 0.10mol, 80% in mineral oil). Stir until solution attained and add 3,4-(dibromomethyl)furan (23.8g, 0.100mol). Stir for 30 minutes then add additional sodium hydride (3.0g, 0.10mol). Stir at room temperature for 20 hours, filter through filter aid and evaporate the solvent in vacuo. Purify by silica gel chromatography to give 5,5-(dicarboethoxy)-2,4,5,6-tetrahydro-cyclopenta[c]furan.
- 20 Dissolve 5,5-(dicarboethoxy)-2,4,5,6-tetrahydro-cyclopenta[c]furan (15.3g, 60.6mmol) in dimethylsulfoxide (140mL). Add water (14mL) and lithium chloride (7.0g, 0.16mol). Heat at reflux for 4 hours, cool and partition between water (150mL) and methylene chloride (2X150mL). Wash the organic phase with water (150mL), dry (MgSO<sub>4</sub>) and pass through a silica gel plug to give 5-(carboethoxy)-2,4,5,6-tetrahydro-cyclopenta[c]furan.
- 25 Dissolve 5-(carboethoxy)-2,4,5,6-tetrahydro-cyclopenta[c]furan (6.2g, 34.1mmol) in ethanol (95%, 150mL) and water (75mL). Add potassium hydroxide (9.5g, 0.17mol) and stir at room temperature for 1 hour. Partition between water (150mL) and ethyl ether (2X150mL). Acidify the aqueous phase with hydrochloric acid to pH 1. Extract with methylene chloride (2X150mL), dry (Na<sub>2</sub>SO<sub>4</sub>) and evaporate the solvent in vacuo to give 5-(carboxy)-2,4,5,6-tetrahydro-cyclopenta[c]furan.
- 30 Dissolve 5-(carboxy)-2,4,5,6-tetrahydro-cyclopenta[c]furan (3.6g, 23.5mmol) in methanol (60mL) and treat with dimethoxypropane (5.8mL, 47mmol) and sulfuric acid (0.8mL). Stir at room temperature for 1 day. Evaporate the solvent in vacuo, dilute with methylene chloride (75mL) and wash with saturated sodium hydrogen carbonate (35mL). Extract the aqueous phase with methylene chloride (30mL), wash combined organics with brine (30mL) and dry (Na<sub>2</sub>SO<sub>4</sub>). Evaporate the solvent in vacuo and pass through a plug of silica gel to give 5-(carbomethoxy)-2,4,5,6-tetrahydro-cyclopenta[c]furan.
- 35 Mix 4-methoxybenzylthiol (3.0g, 19mmol) in sodium hydroxide (20mL of a 2.5N aqueous solution) and methanol (10mL). Add saturated aqueous copper sulfate solution (1.5mL) and stir at room temperature for 2 hours, blowing air over the top of the mixture. Filter, wash solid with water and dry to give 4-methoxybenzyl disulfide as a pale yellow powder (2.71g, 91%).
- 40 Cool lithium hexamethyldisilazane (4.2mL, 4.2mol, 1.0M in tetrahydrofuran) to -78 °C and treat with a solution of 5-(carbomethoxy)-2,4,5,6-tetrahydro-cyclopenta[c]furan (589mg, 3.55mmol) in tetrahydrofuran (5mL). Stir for 1 hour then add hexamethylphosphoramide (0.93mL, 5.3mmol) and stir for 5 minutes. Add 4-methoxybenzyl disulfide (1.6g, 5.2mmol) in tetrahydrofuran (10mL). Stir for 5 hours at -78 °C and quench with a solution of ammonium chloride. Partition between ethyl acetate (75mL) and brine (30mL). Dry (Na<sub>2</sub>SO<sub>4</sub>), evaporate the solvent in vacuo and purify by silica gel chromatography to give 5-(carbomethoxy)-5-(4-methoxybenzylthio)-2,4,5,6-tetrahydro-cyclopenta[c]furan.
- 45 Dissolve 5-(carbomethoxy)-5-(4-methoxybenzylthio)-2,4,5,6-tetrahydro-cyclopenta[c]furan (1.08g, 3.55mmol) in 95% ethanol (25mL), water (12mL) and tetrahydrofuran (15mL).
- 50 Treat with potassium hydroxide (1.3g, 23mmol) and stir at room temperature for 1 hour. Filter and evaporate the solvent in vacuo. Partition between water (125mL) and ether (75mL). Separate the aqueous phase and acidify with cold concentrated hydrochloric acid. Extract with methylene chloride (75mL), dry (Na<sub>2</sub>SO<sub>4</sub>) and evaporate the solvent in vacuo. Purify by silica gel chromatography to give 5-(carboxy)-5-(4-methoxybenzylthio)-2,4,5,6-tetrahydro-cyclopenta[c]furan.
- 55 Dissolve [4S-[4 $\alpha$ , 7 $\alpha$ (R'), 12b $\beta$ ]]-7-(amino)-3,4,6,7,8,12b-hexahydro-6-oxo-1H-[1,4]-thiazino[3,4-a][2]-benzazepine-4-carboxylic acid, diphenylmethyl ester (208mg, 0.454mmol) and EDC (130mg, 0.68mmol) in tetrahydrofuran (5mL). Treat with 5-(carboxy)-5-(4-methoxybenzylthio)-2,4,5,6-tetrahydro-cyclopenta[c]furan (132mg, 0.454mmol). Stir at room temperature for 20 hours and evaporate the solvent in vacuo. Dissolve

the residue in ethyl acetate (50mL) and wash with 5% sulfuric acid (15mL) then saturated sodium hydrogen carbonate (15mL). Dry ( $\text{Na}_2\text{SO}_4$ ) and evaporate the solvent in *vacuo*. Purify by silica gel chromatography to give [4S-[4 $\alpha$ , 7 $\alpha$ (R), 12b $\beta$ ]-7-[(5-(4-methoxybenzylthio)-5-oxo-2,4,5,6-tetrahydro-cyclopenta[c]furan)-methylamino]-3,4,6,7,8,12b-hexahydro-6-oxo-1H-[1,4]-thiazino[3,4-a][2]benzazepine-4-carboxylic acid, diphenylmethyl ester.

Dissolve [4S-[4 $\alpha$ , 7 $\alpha$ (R), 12b $\beta$ ]-7-[(5-(4-methoxybenzylthio)-5-oxo-2,4,5,6-tetrahydro-cyclopenta[c]furan)-methylamino]-3,4,6,7,8,12b-hexahydro-6-oxo-1H-[1,4]-thiazino[3,4-a][2]benzazepine-4-carboxylic acid, diphenylmethyl ester (119mg, 0.163mmol) in methylene chloride (3mL) and cool to 0°C. Treat with trifluoroacetic acid (1.5mL), anisole (0.19mL, 1.7mmol), and mercuric acetate (65mg, 0.2mmol). Stir at 0°C for 3 hours then bubble hydrogen sulfide gas through the solution for 10 minutes. Filter and wash with methylene chloride. Wash organic phase with water (20mL), dry ( $\text{Na}_2\text{SO}_4$ ) and evaporate the solvent in *vacuo*. Purify by silica gel chromatography to give the title compound.

#### Example 15

Preparation of [4S-[4 $\alpha$ , 7 $\alpha$ (R), 12b $\beta$ ]-7-[(5-Thio-5-oxo-4,5,6-trihydro-cyclopenta[c]thiophene)methylamino]-3,4,6,7,8,12b-hexahydro-6-oxo-1H-[1,4]-thiazino[3,4-a][2]benzazepine-4-carboxylic acid

Dissolve 3,4-(dicarboethoxy)thiophene (2.28g, 10mmol) in anhydrous tetrahydrofuran (30mL) and cool to -20°C. Treat with lithium borohydride (7mL of a 2N solution) and stir under a nitrogen atmosphere for several days. Carefully add water and partition between ethyl acetate and 5% hydrochloric acid. Separate the organic phase, wash with brine and dry ( $\text{MgSO}_4$ ). Evaporate the solvent in *vacuo* and purify by silica gel chromatography to give 3,4-(dihydroxymethyl)thiophene.

Dissolve 3,4-(dihydroxymethyl)thiophene (13.4g, 0.093mol) in 2,4,6-collidine (12.4g) and add a solution of lithium bromide (7.9g, 91mmol) in dimethylformamide (80mL). Cool to -50°C and place under a nitrogen atmosphere. Add trifluoromethanesulfonyl chloride (11.8g) at a rate which keeps the temperature under 0°C. Stir at 0°C for 2 hours and pour onto ice water. Extract with ethyl ether, dry ( $\text{MgSO}_4$ ), evaporate the solvent in *vacuo* and purify by silica gel chromatography to give 3,4-(dibromomethyl)thiophene.

Dissolve diethylmalonate (15.2mL, 0.100mol) in tetrahydrofuran (800mL). Cool in an ice bath and treat with sodium hydride (3.0g, 0.10mol, 80% in mineral oil). Stir until solution attained and add 3,4-(dibromomethyl)thiophene (25.4g, 0.100mol). Stir for 30 minutes then add additional sodium hydride (3.0g, 0.10mol). Stir at room temperature for 20 hours, filter through filter aid and evaporate the solvent in *vacuo*. Purify by silica gel chromatography to give 5,5-(dicarboethoxy)-2,4,5,6-tetrahydro-cyclopenta[c]thiophene.

Dissolve 5,5-(dicarboethoxy)-2,4,5,6-tetrahydro-cyclopenta[c]thiophene (16.2g, 60.6mmol) in dimethylsulfoxide (140mL). Add water (14mL) and lithium chloride (7.0g, 0.16mol). Heat at reflux for 4 hours, cool and partition between water (150mL) and methylene chloride (2X150mL). Wash the organic phase with water (150mL), dry ( $\text{MgSO}_4$ ) and pass through a silica gel plug to give 5-(carboethoxy)-2,4,5,6-tetrahydro-cyclopenta[c]thiophene.

Dissolve 5-(carboethoxy)-2,4,5,6-tetrahydro-cyclopenta[c]thiophene (6.68g, 34.1mmol) in ethanol (95%, 150mL) and water (75mL). Add potassium hydroxide (9.5g, 0.17mol) and stir at room temperature for 1 hour. Partition between water (150mL) and ethyl ether (2X150mL). Acidify the aqueous phase with hydrochloric acid to pH 1. Extract with methylene chloride (2X150mL), dry ( $\text{Na}_2\text{SO}_4$ ) and evaporate the solvent in *vacuo* to give 5-(carboxy)-2,4,5,6-tetrahydro-cyclopenta[c]thiophene.

Dissolve 5-(carboxy)-2,4,5,6-tetrahydro-cyclopenta[c]thiophene (3.95g, 23.5mmol) in methanol (60mL) and treat with dimethoxypropane (5.8mL, 47mmol) and sulfuric acid (0.8mL). Stir at room temperature for 1 day. Evaporate the solvent in *vacuo*, dilute with methylene chloride (75mL) and wash with saturated sodium hydrogen carbonate (35mL). Extract the aqueous phase with methylene chloride (30mL), wash combined organics with brine (30mL) and dry ( $\text{Na}_2\text{SO}_4$ ). Evaporate the solvent in *vacuo* and pass through a plug of silica gel to give 5-(carboethoxy)-2,4,5,6-tetrahydro-cyclopenta[c]thiophene.

Mix 4-methoxybenzylthiol (3.0g, 19mmol) in sodium hydroxide (20mL of a 2.5N aqueous solution) and methanol (10mL). Add saturated aqueous copper sulfate solution (1.5mL) and stir at room temperature for 2 hours, blowing air over the top of the mixture. Filter, wash solid with water and dry to give 4-methoxybenzyl disulfide as a pale yellow powder (2.71g, 91%).

Cool lithium hexamethyldisilazane (4.2mL, 4.2mol, 1.0M in tetrahydrofuran) to -78°C and treat with a solution of 5-(carboethoxy)-2,4,5,6-tetrahydro-cyclopenta[c]thiophene (646mg, 3.55mmol) in tetrahydrofuran (5mL). Stir for 1 hour then add hexamethylphosphoramide (0.93mL, 5.3mmol) and stir for 5 minutes. Add 4-methoxybenzyl disulfide (1.6g, 5.2mmol) in tetrahydrofuran (10mL). Stir for 5 hours at -78°C and quench with a solution of ammonium chloride. Partition between ethyl acetate (75mL) and brine

- (30mL). Dry ( $\text{Na}_2\text{SO}_4$ ), evaporate the solvent in vacuo and purify by silica gel chromatography to give 5-(carbomethoxy)-5-(4-methoxybenzylthio)-2,4,5,6-tetrahydro-cyclopenta[c]thiophene.
- Dissolve 5-(carbomethoxy)-5-(4-methoxybenzylthio)-2,4,5,6-tetrahydro-cyclopenta[c]thiophene (1.14g, 3.55mmol) in 95% ethanol (25mL), water (12mL) and tetrahydrofuran (15mL). Treat with potassium hydroxide (1.3g, 23mmol) and stir at room temperature for 1 hour. Filter and evaporate the solvent in vacuo. Partition between water (125mL) and ether (75mL). Separate the aqueous phase and acidify with cold concentrated hydrochloric acid. Extract with methylene chloride (75mL), dry ( $\text{Na}_2\text{SO}_4$ ) and evaporate the solvent in vacuo. Purify by silica gel chromatography to give 5-(carboxy)-5-(4-methoxybenzylthio)-2,4,5,6-tetrahydro-cyclopenta[c]thiophene.
- Dissolve [4S-[4 $\alpha$ , 7 $\alpha$ (R\*), 12b $\beta$ ]]-7-(amino)-3,4,6,7,8,12b-hexahydro-6-oxo-1H-[1,4]-thiazino[3,4-a][2]-benzazepine-4-carboxylic acid, diphenylmethyl ester (208mg, 0.454mmol) and EDC (130mg, 0.68mmol) in tetrahydrofuran (5mL). Treat with 5-(carboxy)-5-(4-methoxybenzylthio)-2,4,5,6-tetrahydro-cyclopenta[c]thiophene (139mg, 0.454mmol). Stir at room temperature for 20 hours and evaporate the solvent in vacuo. Dissolve the residue in ethyl acetate (50mL) and wash with 5% sulfuric acid (15mL) then saturated sodium hydrogen carbonate (15mL). Dry ( $\text{Na}_2\text{SO}_4$ ) and evaporate the solvent in vacuo. Purify by silica gel chromatography to give [4S-[4 $\alpha$ , 7 $\alpha$ (R\*), 12b $\beta$ ]]-7-[(5-(4-methoxybenzylthio)-5-oxo-2,4,5,6-tetrahydro-cyclopenta[c]thiophene)methylamino]-3,4,6,7,8,12b-hexahydro-6-oxo-1H-[1,4]-thiazino[3,4-a][2]benzazepine-4-carboxylic acid, diphenylmethyl ester.
- Dissolve [4S-[4 $\alpha$ , 7 $\alpha$ (R\*), 12b $\beta$ ]]-7-[(5-(4-methoxybenzylthio)-5-oxo-2,4,5,6-tetrahydro-cyclopenta[c]thiophene)methylamino]-3,4,6,7,8,12b-hexahydro-6-oxo-1H-[1,4]-thiazino[3,4-a][2]benzazepine-4-carboxylic acid, diphenylmethyl ester (122mg, 0.163mmol) in methylene chloride (3mL) and cool to 0 °C. Treat with trifluoroacetic acid (1.5mL), anisole (0.19mL, 1.7mmol), and mercuric acetate (65mg, 0.2mmol). Stir at 0 °C for 3 hours then bubble hydrogen sulfide gas through the solution for 10 minutes. Filter and wash with methylene chloride. Wash organic phase with water (20mL), dry ( $\text{Na}_2\text{SO}_4$ ) and evaporate the solvent in vacuo. Purify by silica gel chromatography to give the title compound.

#### Example 16

- Preparation of [4S-[4 $\alpha$ , 7 $\alpha$ (R\*), 12b $\beta$ ]]-7-[(5-Thio-5-oxo-2,4,5,6-tetrahydro-cyclopenta[c]pyrrole)methylamino]-3,4,6,7,8,12b-hexahydro-6-oxo-1H-[1,4]-azazino[3,4-a][2]benzazepine-4-carboxylic acid

Scheme E, step a: N-[2-(1,3-Dihydro-1,3-dioxo-2H-isoindol-2-yl)-1-oxo-3-phenylpropyl]-(S)-3-[(trifluoroacetyl-2-propenyl)amino]-2-amino-propionic acid, methyl ester

- Dissolve N<sup>+</sup>-(carbobenzyloxy)- $\beta$ -(amino)-L-alanine (47.6g, 0.2mol) in methanol (500mL) and treat with concentrated sulfuric acid (0.5mL). Heat to 60 °C for 16 hours, cool and reduce the solvent by 50% in vacuo. Dilute with ethyl ether (500mL), wash with saturated sodium hydrogen carbonate, then brine. Dry ( $\text{MgSO}_4$ ) and evaporate the solvent in vacuo to give N<sup>+</sup>-(carbobenzyloxy)- $\beta$ -(amino)-L-alanine, methyl ester.
- Dissolve N<sup>+</sup>-(carbobenzyloxy)- $\beta$ -(amino)-L-alanine, methyl ester (15.9g, 63mmol) in methylene chloride/cyclohexane (1:1, 600mL). Add allyl trichloroacetimidate (26g, 128mmol) and trifluoromethanesulfonic acid (5mL, 56.6mmol). Stir at room temperature under a nitrogen atmosphere for 5 hours and dilute with methylene chloride. Wash with saturated aqueous sodium hydrogen carbonate, water, dry ( $\text{MgSO}_4$ ) and evaporate the solvent in vacuo. Purify by silica gel chromatography to give N<sup>+</sup>-(carbobenzyloxy)- $\beta$ -(allylamino)-L-alanine, methyl ester.
- Dissolve N<sup>+</sup>-(carbobenzyloxy)- $\beta$ -(allylamino)-L-alanine, methyl ester (663mg, 2.27mmol) in anhydrous tetrahydrofuran (15mL). Treat with pyridine (183 $\mu$ L, 2.27mmol) followed by trifluoroacetic anhydride (321 $\mu$ L, 2.27mmol) and stir at room temperature overnight. Partition between ethyl ether and water. Separate the organic phase, dry ( $\text{MgSO}_4$ ) and evaporate the solvent in vacuo. Purify by silica gel chromatography to give N<sup>+</sup>-(carbobenzyloxy)- $\beta$ -(trifluoroacetyl-allylamino)-L-alanine, methyl ester.
- Place boron tribromide (215mg, 0.86mmol) in a flask and cool to 0 °C. Cautiously add trifluoroacetic acid (5mL) with stirring. Evaporate the solvent to give boron tris(trifluoroacetate).
- Dissolve boron tris(trifluoroacetate) (0.3g, 0.86mmol) in trifluoroacetic acid (10mL) and add N<sup>+</sup>-(carbobenzyloxy)- $\beta$ -(trifluoroacetyl-allylamino)-L-alanine, methyl ester (105mg, 0.27mmol). Stir under an argon atmosphere for 1 hour then evaporate the solvent in vacuo at room temperature. Add methanol and evaporate repeatedly. Purify by silica gel chromatography to give  $\beta$ -(trifluoroacetyl-allylamino)-L-alanine, methyl ester, hydrochloride.
- Dissolve  $\beta$ -(trifluoroacetyl-allylamino)-L-alanine, methyl ester, hydrochloride (104.8g, 0.36mol) in tetrahydrofuran (300mL) then cool to 0 °C and add 4-methylmorpholine (88mL, 0.8mol). Add, by dropwise

addition, a solution of the N-phthaloyl-(S)-phenylalanine, acid chloride (108.7g, 0.36mol) in tetrahydrofuran (200mL). Allow to warm to room temperature and stir for 3 hours. Filter and concentrate the filtrate in vacuo. Dissolve the residue in ethyl acetate and separate the organic phase. Wash with water then saturated sodium chloride and dry (MgSO<sub>4</sub>). Evaporate the solvent in vacuo to give an oil. Purify by silica gel chromatography to give the title compound.

Scheme E, step b: [S-(R\*, R\*)]-N-[2-(1,3-Dihydro-1,3-dioxo-2H-isoindol-2-yl)-1-oxo-3-phenylpropyl]-3,4-dihydro-2H-4-trifluoroacetyl-1,4-azazine-3-carboxylic acid, methyl ester Dissolve N-[2-(1,3-dihydro-1,3-dioxo-2H-isoindol-2-yl)-1-oxo-3-phenylpropyl]-(S)-3-[(trifluoroacetyl-2-propenyl)amino]-2-amino-propionic acid, methyl ester (15.8g, 29.8mmol) in methylene chloride/methanol (10:1, 220mL). Cool to -78 °C and sparge with a mixture of ozone/oxygen for approximately 10 minutes until a blue color persists. Sparge with nitrogen for 10 minutes at -78 °C to remove excess ozone. Treat with methylsulfide (60mL, 0.82mol) and allow to warm to room temperature. Stir at room temperature for 2.5 hours, evaporate the solvent in vacuo and dissolve the residue in ethyl acetate (200mL). Wash with water, saturated sodium chloride, dry (MgSO<sub>4</sub>) and evaporate the solvent in vacuo to give the intermediate N-[2-(1,3-dihydro-1,3-dioxo-2H-isoindol-2-yl)-1-oxo-3-phenylpropyl]-N-2-oxoethyl, methyl ester. Dissolve N-[2-(1,3-dihydro-1,3-dioxo-2H-isoindol-2-yl)-1-oxo-3-phenylpropyl]-(S)-3-[(trifluoroacetyl-2-oxoethyl)amino]-2-amino-propionic acid, methyl ester (15.9g, 29.8mmol) in methylene chloride/trifluoroacetic acid (10:1/330mL). Stir at room temperature for 2.5 hours and evaporate the solvent in vacuo. Purify by silica gel chromatography to give the title compound.

Scheme E, step c: [4S-[4 $\alpha$ , 7 $\alpha$ (R\*), 12b $\beta$ ]]-7-[(1,3-Dihydro-1,3-dioxo-2H-isoindol-2-yl)]-3,4,6,7,8,12b-hexahydro-6-oxo-1H-4-trifluoroacetyl-[1,4]-azazino[3,4-a][2]benzazepine-4-carboxylic acid, diphenylmethyl ester

Dissolve [S-(R\*, R\*)]-N-[2-(1,3-dihydro-1,3-dioxo-2H-isoindol-2-yl)-1-oxo-3-phenylpropyl]-3,4-dihydro-2H-4-trifluoroacetyl-1,4-azazine-3-carboxylic acid, methyl ester (3.04g, 5.9mmol) in methylene chloride (5mL) and add, by dropwise addition, to a previously prepared solution of trifluoromethanesulfonic acid (4.0mL, 45mmol) and trifluoroacetic anhydride (1.0mL, 7.1mmol). Place under a nitrogen atmosphere and stir at room temperature for 123 hours. Pour into a separatory funnel containing ice (200g) and ethyl acetate (200mL). Separate the organic phase, wash with water (3X200mL) and saturated aqueous sodium chloride (100mL). Extract the organic phase with 10% wt. potassium hydrogen carbonate (4X40mL) and water (40mL). Layer the combined basic aqueous phases with ethyl acetate (100mL) and cool in an ice bath. Add, by dropwise addition, 6N hydrochloric acid to adjust the pH to 1 while maintaining the temperature at 5-10 °C. Separate the organic phase and extract the aqueous phase with ethyl acetate (3X200mL), wash with saturated sodium chloride and dry (MgSO<sub>4</sub>). Evaporate the solvent in vacuo and dry the residue under high vacuum at 56 °C for 24 hours to give the intermediate [4S-[4 $\alpha$ , 7 $\alpha$ (R\*), 12b $\beta$ ]]-7-[(1,3-dihydro-1,3-dioxo-2H-isoindol-2-yl)]-3,4,6,7,8,12b-hexahydro-6-oxo-1H-4-trifluoroacetyl-[1,4]-oxazino[3,4-a][2]benzazepine-4-carboxylic acid. Dissolve [4S-[4 $\alpha$ , 7 $\alpha$ (R\*), 12b $\beta$ ]]-7-[(1,3-dihydro-1,3-dioxo-2H-isoindol-2-yl)]-3,4,6,7,8,12b-hexahydro-6-oxo-1H-4-trifluoroacetyl-[1,4]-azazino[3,4-a][2]benzazepine-4-carboxylic acid (616mg, 1.23mmol) in methylene chloride (12mL) and treat with diphenyldiazomethane (360mg, 1.86mmol). Stir for 5.5 hours and evaporate the solvent in vacuo. Purify by silica gel chromatography to give the title compound.

Scheme E, step e: [4S-[4 $\alpha$ , 7 $\alpha$ (R\*), 12b $\beta$ ]]-7-(Amino)-3,4,6,7,8,12b-hexahydro-6-oxo-1H-[1,4]-azazino[3,4-a][2]benzazepine-4-carboxylic acid, diphenylmethyl ester

Dissolve [4S-[4 $\alpha$ , 7 $\alpha$ (R\*), 12b $\beta$ ]]-7-[(1,3-dihydro-1,3-dioxo-2H-isoindol-2-yl)]-3,4,6,7,8,12b-hexahydro-6-oxo-1H-4-trifluoroacetyl-[1,4]-azazino[3,4-a][2]benzazepine-4-carboxylic acid, diphenylmethyl ester (345mg, 0.517mmol) in methanol (5mL) and treat with hydrazine monohydrate (1.1mL of a 1M solution in methanol, 1.1mmol). Stir at room temperature for 44 hours, evaporate the solvent in vacuo and slurry the residue in methylene chloride (10mL). Filter and evaporate the solvent in vacuo to give the title compound.

Scheme A: [4S-[4 $\alpha$ , 7 $\alpha$ (R\*), 12b $\beta$ ]]-7-[(5-Thio-5-oxo-2,4,5,6-tetrahydro-cyclopenta[c]pyrrole)methylamino]-3,4,6,7,8,12b-hexahydro-6-oxo-1H-[1,4]-azazino[3,4-a][2]benzazepine-4-carboxylic acid

Dissolve 3,4-(dicarboethoxy)pyrrole (1.06g, 5mmol) in 50/50 dioxane water (25mL) and buffer to pH 10 with 1N sodium hydroxide. Add, by dropwise addition, an ether solution of di-t-butyl dicarbonate (1.2g, 5.5mmol)

- at 10° C. Allow to warm to room temperature and buffer occasionally to retain pH 10. Acidify with a sodium citrate/citric acid buffer to pH 5, extract with ethyl ether (3X), dry (MgSO<sub>4</sub>) and evaporate the solvent in vacuo. Purify the residue by silica gel chromatography to give N-(carbo-t-butyloxy)-3,4-(dicarboethoxy)-pyrrole.
- 5 Dissolve N-(carbo-t-butyloxy)-3,4-(dicarboethoxy)pyrrole (3.11g, 10mmol) in anhydrous tetrahydrofuran (30mL) and cool to -20° C. Treat with lithium borohydride (7mL of a 2N solution) and stir under a nitrogen atmosphere for several days. Carefully add water and partition between ethyl acetate and 5% hydrochloric acid. Separate the organic phase, wash with brine and dry (MgSO<sub>4</sub>). Evaporate the solvent in vacuo and purify by silica gel chromatography to give N-(carbo-t-butyloxy)-3,4-(dihydroxymethyl)pyrrole.
- 10 Dissolve N-bromosuccinimide (1.78g, 0.01mol) in tetrahydrofuran (60mL) and add a solution of triphenylphosphine (2.62g, 0.01mol) in tetrahydrofuran. Add a solution of N-(carbo-t-butyloxy)-3,4-(dihydroxymethyl)pyrrole (1.14g, 5mmol) in tetrahydrofuran (25mL) and stir until most of the solid goes into solution. Evaporate the solvent in vacuo and partition the residue between water and ethyl ether. Separate the organic phase and wash with water. Dry (Na<sub>2</sub>SO<sub>4</sub>) and evaporate the solvent in vacuo. Purify by silica gel chromatography to give N-(carbo-t-butyloxy)-3,4-(dibromomethyl)pyrrole.
- 15 Dissolve diethylmalonate (15.2mL, 0.100mol) in tetrahydrofuran (800mL). Cool in an ice bath and treat with sodium hydride (3.0g, 0.10mol, 80% in mineral oil). Stir until solution attained and add N-(carbo-t-butyloxy)-3,4-(dibromomethyl)pyrrole (35.3g, 0.100mol). Stir for 30 minutes then add additional sodium hydride (3.0g, 0.10mol). Stir at room temperature for 20 hours, filter through filter aid and evaporate the solvent in vacuo.
- 20 Purify by silica gel chromatography to give 5,5-(dicarboethoxy)-2-(carbo-t-butyloxy)-2,4,5,6-tetrahydro-cyclopenta[c]pyrrole.  
Dissolve 5,5-(dicarboethoxy)-2-(carbo-t-butyloxy)-2,4,5,6-tetrahydro-cyclopenta[c]pyrrole (21.3g, 60.6mmol) in dimethylsulfoxide (140mL). Add water (14mL) and lithium chloride (7.0g, 0.16mol). Heat at reflux for 4 hours, cool and partition between water (150mL) and methylene chloride (2X150mL). Wash the organic phase with water (150mL), dry (MgSO<sub>4</sub>) and pass through a silica gel plug to give 5-(carboethoxy)-2-(carbo-t-butyloxy)-2,4,5,6-tetrahydro-cyclopenta[c]pyrrole.
- 25 Dissolve 5-(carboethoxy)-2-(carbo-t-butyloxy)-2,4,5,6-tetrahydro-cyclopenta[c]pyrrole (9.5g, 34.1mmol) in ethanol (95%, 150mL) and water (75mL). Add potassium hydroxide (9.5g, 0.17mol) and stir at room temperature for 1 hour. Partition between water (150mL) and ethyl ether (2X150mL). Acidify the aqueous phase with hydrochloric acid to pH 1. Extract with methylene chloride (2X150mL), dry (Na<sub>2</sub>SO<sub>4</sub>) and evaporate the solvent in vacuo to give 5-(carboxy)-2-(carbo-t-butyloxy)-2,4,5,6-tetrahydro-cyclopenta[c]pyrrole.
- 30 Dissolve 5-(carboxy)-2-(carbo-t-butyloxy)-2,4,5,6-tetrahydro-cyclopenta[c]pyrrole (5.9g, 23.5mmol) in methanol (60mL) and treat with dimethoxypropane (5.8mL, 47mmol) and sulfuric acid (0.8mL). Stir at room temperature for 1 day. Evaporate the solvent in vacuo, dilute with methylene chloride (75mL) and wash with saturated sodium hydrogen carbonate (35mL). Extract the aqueous phase with methylene chloride (30mL), wash combined organics with brine (30mL) and dry (Na<sub>2</sub>SO<sub>4</sub>). Evaporate the solvent in vacuo and pass through a plug of silica gel to give 5-(carbomethoxy)-2-(carbo-t-butyloxy)-2,4,5,6-tetrahydro-cyclopenta[c]pyrrole.
- 35 Mix 4-methoxybenzylthiol (3.0g, 19mmol) in sodium hydroxide (20mL of a 2.5N aqueous solution) and methanol (10mL). Add saturated aqueous copper sulfate solution (1.5mL) and stir at room temperature for 2 hours, blowing air over the top of the mixture. Filter, wash solid with water and dry to give 4-methoxybenzyl disulfide as a pale yellow powder (2.71g, 91%).
- Cool lithium hexamethyldisilazane (4.2mL, 4.2mol, 1.0M in tetrahydrofuran) to -78° C and treat with a solution of 5-(carbomethoxy)-2-(carbo-t-butyloxy)-2,4,5,6-tetrahydro-cyclopenta[c]pyrrole (941mg, 3.55mmol) in tetrahydrofuran (5mL). Stir for 1 hour then add hexamethylphosphoramide (0.93mL, 5.3mmol) and stir for 5 minutes. Add 4-methoxybenzyl disulfide (1.6g, 5.2mmol) in tetrahydrofuran (10mL). Stir for 5 hours at -78° C and quench with a solution of ammonium chloride. Partition between ethyl acetate (75mL) and brine (30mL). Dry (Na<sub>2</sub>SO<sub>4</sub>), evaporate the solvent in vacuo and purify by silica gel chromatography to give 5-(carbomethoxy)-2-(carbo-t-butyloxy)-5-(4-methoxybenzylthio)-2,4,5,6-tetrahydro-cyclopenta[c]pyrrole.
- 50 Dissolve 5-(carbomethoxy)-2-(carbo-t-butyloxy)-5-(4-methoxybenzylthio)-2,4,5,6-tetrahydro-cyclopenta[c]pyrrole (1.43g, 3.55mmol) in 95% ethanol (25mL), water (12mL) and tetrahydrofuran (15mL). Treat with potassium hydroxide (1.3g, 23mmol) and stir at room temperature for 1 hour. Filter and evaporate the solvent in vacuo. Partition between water (125mL) and ether (75mL). Separate the aqueous phase and acidify with cold concentrated hydrochloric acid. Extract with methylene chloride (75mL), dry (Na<sub>2</sub>SO<sub>4</sub>) and evaporate the solvent in vacuo. Purify by silica gel chromatography to give 5-(carboxy)-2-(carbo-t-butyloxy)-5-(4-methoxybenzylthio)-2,4,5,6-tetrahydro-cyclopenta[c]pyrrole.
- 55 Dissolve [4S-[4 $\alpha$ , 7 $\alpha$ (R), 12b $\beta$ ]]-7-(amino)-3,4,6,7,8,12b-hexahydro-6-oxo-1H-[1,4]-azazino[3,4-a][2-

benzazepine-4-carboxylic acid, diphenylmethyl ester (200mg, 0.454mmol) and EDC (130mg, 0.68mmol) in tetrahydrofuran (5mL). Treat with 5-(carboxy)-2-(carbo-t-butyloxy)-5-(4-methoxybenzylthio)-2,4,5,6-tetrahydro-cyclopenta[c]pyrrole (177mg, 0.454mmol). Stir at room temperature for 20 hours and evaporate the solvent in vacuo. Dissolve the residue in ethyl acetate (50mL) and wash with 5% sulfuric acid (15ml) then saturated sodium hydrogen carbonate (15mL). Dry (Na<sub>2</sub>SO<sub>4</sub>) and evaporate the solvent in vacuo. Purify by silica gel chromatography to give [4S-[4 $\alpha$ , 7 $\alpha$ (R\*), 12b $\beta$ ]]-7-[(5-(4-methoxybenzylthio)-5-oxo-2-(carbo-t-butyloxy)-2,4,5,6-tetrahydro-cyclopenta[c]pyrrole)methylamino]-3,4,6,7,8,12b-hexahydro-6-oxo-1H-[1,4]-azazino[3,4-a]-[2]benzazepine-4-carboxylic acid, diphenylmethyl ester.

Dissolve [4S-[4 $\alpha$ , 7 $\alpha$ (R\*), 12b $\beta$ ]]-7-[(5-(4-methoxybenzylthio)-5-oxo-2-(carbo-t-butyloxy)-2,4,5,6-tetrahydro-cyclopenta[c]pyrrole)methylamino]-3,4,6,7,8,12b-hexahydro-6-oxo-1H-[1,4]-azazino[3,4-a][2]benzazepine-4-carboxylic acid, diphenylmethyl ester (132mg, 0.163mmol) in methylene chloride (3mL) and cool to 0 °C. Treat with trifluoroacetic acid (1.5mL), anisole (0.19mL, 1.7mmol), and mercuric acetate (65mg, 0.2mmol). Stir at 0 °C for 3 hours then bubble hydrogen sulfide gas through the solution for 10 minutes. Filter and wash with methylene chloride. Wash organic phase with water (20mL), dry (Na<sub>2</sub>SO<sub>4</sub>) and evaporate the solvent in vacuo. Purify by silica gel chromatography to give the title compound.

#### Example 17

Preparation of [4S-[4 $\alpha$ , 7 $\alpha$ (R\*), 12b $\beta$ ]]-7-1(5-Thio-5-oxo-2,4,5,6-tetrahydro-cyclopenta[c]pyrrole)-methylamino]-3,4,6,7,8,12b-hexahydro-6-oxo-1H-[1,4]-N<sup>4</sup>-trifluoroacetyl-azazino[3,4-a][2]benzazepine-4-carboxylic acid

Dissolve [4S-[4 $\alpha$ , 7 $\alpha$ (R\*), 12b $\beta$ ]]-7-[(5-(4-methoxybenzylthio)-5-oxo-2-(carbo-t-butyloxy)-2,4,5,6-tetrahydro-cyclopenta[c]pyrrole)methylamino]-3,4,6,7,8,12b-hexahydro-6-oxo-1H-[1,4]-azazino[3,4-a][2]benzazepine-4-carboxylic acid, diphenylmethyl ester (1.84g, 2.27mmol) in anhydrous tetrahydrofuran (15mL). Treat with pyridine (183 $\mu$ L, 2.27mmol) followed by trifluoroacetic anhydride (321 $\mu$ L, 2.27mmol) and stir at room temperature overnight. Partition between ethyl ether and water. Separate the organic phase, dry (MgSO<sub>4</sub>) and evaporate the solvent in vacuo. Purify by silica gel chromatography to give [4S-[4 $\alpha$ , 7 $\alpha$ (R\*), 12b $\beta$ ]]-7-[(5-(4-methoxybenzylthio)-5-oxo-2-(carbo-t-butyloxy)-2,4,5,6-tetrahydro-cyclopenta[c]pyrrole)methylamino]-3,4,6,7,8,12b-hexahydro-6-oxo-1H-[1,4]-N<sup>4</sup>-trifluoroacetyl-azazino[3,4-a][2]benzazepine-4-carboxylic acid, diphenylmethyl ester.

Dissolve [4S-[4 $\alpha$ , 7 $\alpha$ (R\*), 12b $\beta$ ]]-7-[(5-(4-methoxybenzylthio)-5-oxo-2-(carbo-t-butyloxy)-2,4,5,6-tetrahydro-cyclopenta[c]pyrrole)methylamino]-3,4,6,7,8,12b-hexahydro-6-oxo-1H-[1,4]-N<sup>4</sup>-trifluoroacetyl-azazino[3,4-a][2]benzazepine-4-carboxylic acid, diphenylmethyl ester (148mg, 0.163mmol) in methylene chloride (3mL) and cool to 0 °C. Treat with trifluoroacetic acid (1.5mL), anisole (0.19mL, 1.7mmol), and mercuric acetate (65mg, 0.2mmol). Stir at 0 °C for 3 hours then bubble hydrogen sulfide gas through the solution for 10 minutes. Filter and wash with methylene chloride. Wash organic phase with water (20mL), dry (Na<sub>2</sub>SO<sub>4</sub>) and evaporate the solvent in vacuo. Purify by silica gel chromatography to give the title compound.

#### Example 18

[6 $\alpha$ (R\*), 11b $\beta$ ]-6-[(S)-(5-Thio-5-oxo-2,4,5,6-tetrahydro-cyclopenta[c]pyrrole)methylamino]-1,2,3,5,6,7,11b-heptahydro-5-oxo-pyrrolo[2,1-a][2]benzazepine-3(S)-carboxylic acid, methyl ester

#### 45 Scheme G, step a: N-(Phenylmethylene)-2-(3-butenyl)glycine methyl ester

Dissolve diisopropylamine (15.4mL, 110mmol) in tetrahydrofuran (250mL), place under a nitrogen atmosphere and cool to -78 °C. Add n-butyllithium (39mL of a 2.7M solution in hexane, 105mmol). Stir for 30 minutes and add, by dropwise addition, a solution of N-(phenylmethylene)glycine methyl ester (17.7g, 100mmol) in tetrahydrofuran (25mL). Stir for 15 minutes and add 4-bromobutene (13.5g, 100mmol) and allow to warm slowly to room temperature. Add hexamethylphosphoramide (20mL, 100mmol) and stir under a nitrogen atmosphere for 3 hours.

Pour into water, extract into ethyl ether and wash with brine several times. Dry (MgSO<sub>4</sub>) and evaporate the solvent in vacuo to give the title compound as an amber oil (25g).

45 Scheme G, step b: 2-(3-Butenyl)glycine methyl ester Dissolve N-(phenylmethylene)-2-(3-butenyl)glycine methyl ester (25g) in ethyl ether (400mL) and stir with 1N hydrochloric acid (150mL) and water (150mL). Place under an argon atmosphere and stir for 2 hours. Separate the aqueous phase and adjust to pH 9, extract into chloroform, dry and evaporate the solvent in vacuo to give the title compound as a light oil



(4.5g).

Scheme G, step c: (S)-N-[2-(1,3-Dihydro-1,3-dioxo-2H-isoindol-2-yl)-1-oxo-3-phenylpropyl]-2-(3-butenyl)-glycine, methyl esters

5 Dissolve N-phthaloyl-(S)-phenylalanine (2) (6.0g, 20mmol) and EEDQ (6.0g, 24mmol) in methylene chloride (30mL). Add 2-(3-butenyl)glycine methyl ester (3.0g, 21mmol) and stir for 18 hours. Pour into methylene chloride, wash with 10% hydrochloric acid (2X100mL) then saturated sodium hydrogen carbonate. Dry and evaporate the solvent in vacuo to give 8.3g yellow oil. Purify by silica gel chromatography (25% ethyl acetate/hexane) to give a diastereomeric mixture of the title compounds as foam (5.2g).

Scheme G, step d: (S)-N-[2-(1,3-Dihydro-1,3-dioxo-2H-isoindol-2-yl)]-phenylalanyl]-2-(3-oxopropyl)glycine, methyl esters

15 Dissolve the diastereomeric mixture of (S)-N-[2-(1,3-Dihydro-1,3-dioxo-2H-isoindol-2-yl)-1-oxo-3-phenylpropyl]-2-(3-butenyl)-glycine, methyl esters (4.2g, 10mmol) in methylene chloride (100mL) and absolute methanol (10mL). Cool to -78 °C and treat with ozone until blue. Degas with oxygen and add methyl sulfide (10mL) and pyridine (0.5mL). Allow to warm slowly to room temperature and stir for 18 hours. Wash with 10% hydrochloric acid then brine. Dry and evaporate the solvent in vacuo to give a diastereomeric mixture of the title compounds as an oil (4.5g).

Scheme F, step c: (S)-N-[2-(1,3-Dihydro-1,3-dioxo-2H-isoindol-2-yl)]-1-oxo-3-phenylpropyl-1,2,3-trihydro-2-(S)-pyrrolicarboxylic acid, methyl ester and (S)-N-[2-(1,3-Dihydro-1,3-dioxo-2H-isoindol-2-yl)]-1-oxo-3-phenylpropyl-1,2,3-trihydro-2(R)-pyrrolicarboxylic acid, methyl ester

25 Dissolve the diastereomeric mixture of (S)-N-[2-(1,3-Dihydro-1,3-dioxo-2H-isoindol-2-yl)]-(S)-phenylalanyl]-2-(3-oxopropyl)glycine, methyl esters (4.5g) in 1,1,1-trichloroethane (150mL) and treat with trifluoroacetic acid (0.5mL). Heat at reflux for 18 hours, evaporate the solvent and purify by silica gel chromatography (80% ethyl acetate/hexane) to give the 2(S)-title compound (700mg) and the 2(R)-title compound (600mg).

Scheme F, Step d: [6 $\alpha$ (R\*), 11b $\beta$ ]-6-[(S)-(1,3-Dihydro-1,3-dioxo-2H-isoindol-2-yl)]-1,2,3,5,6,7,11b-heptahydro-5-oxo-pyrrolo[2,1-a][2]benzazepine-3(S)-carboxylic acid, methyl ester

35 Dissolve (S)-N-[2-(1,3-dihydro-1,3-dioxo-2H-isoindol-2-yl)]-1-oxo-3-phenylpropyl-1,2,3-trihydro-2(S)-pyrrolicarboxylic acid, methyl ester (338mg, 0.836mmol) in anhydrous methylene chloride (10mL) and add to trifluoromethanesulfonic acid (5mL). Stir for 3.5 hours, cool in an ice bath and carefully add water (25mL). Extract with ethyl acetate (75mL) and wash with saturated sodium hydrogen carbonate (25mL). Dry (Na<sub>2</sub>SO<sub>4</sub>) and evaporate the solvent in vacuo. Purify by silica gel chromatography (1:1 ethyl acetate/hexane to 2:1 ethyl acetate/hexane) to give the title compound as a white foam (314mg, 93%).

Scheme F, Step e: [6 $\alpha$ (R\*), 11b $\beta$ ]-6-[(S)-Amino]-1,2,3,5,6,7,11b-heptahydro-5-oxo-pyrrolo[2,1-a][2]benzazepine-3(S)-carboxylic acid, methyl ester

45 Dissolve [6 $\alpha$ (R\*), 11b $\beta$ ]-6-[(S)-(1,3-Dihydro-1,3-dioxo-2H-isoindol-2-yl)]-1,2,3,5,6,7,11b-heptahydro-5-oxo-pyrrolo[2,1-a][2]benzazepine-3(S)-carboxylic acid, methyl ester (244mg, 0.603mmol) in methanol (3mL) and treat with hydrazine monohydrate (0.70mL of a 1M solution in methanol) and stir at room temperature for 24 hours. Add additional hydrazine monohydrate (0.3mL of a 1M solution in methanol) and stir for 48 hours. Filter through filter aid, evaporate the solvent in vacuo and add methylene chloride. Filter slowly through filter aid and evaporate the solvent in vacuo to give the title compound as a yellow oil (181mg).

Scheme A: [6 $\alpha$ (R\*), 11b $\beta$ ]-6-[(S)-(5-Thio-5-oxo-2,4,5,6-tetrahydro-cyclopenta[c]pyrrole)methylamino]-1,2,3,5,6,7,11b-heptahydro-5-oxo-pyrrolo[2,1-a][2]benzazepine-3(S)-carboxylic acid, methyl ester

55 Dissolve 5-(carboxy)-2-(carbo-t-butyloxy)-5-(4-methoxybenzylthio)-2,4,5,6-tetrahydro-cyclopenta[c]pyrrole (329mg, 0.845mmol) in methylene chloride (6mL), cool in an ice-methanol bath and treat with oxalyl chloride (0.94mL, 11mmol). Stir for 1.5 hours, evaporate the solvent in vacuo at 0-5 °C. Dilute the residue with methylene chloride (3mL) and add a solution of [6 $\alpha$ (R\*), 11b $\beta$ ]-6-[(S)-Amino]-1,2,3,5,6,7,11b-heptahydro-5-oxo-pyrrolo[2,1-a][2]benzazepine-3(S)-carboxylic acid, methyl ester (155mg, 0.565mmol) in

methylene chloride (6mL). Add pyridine (68μL, 0.85mmol) and stir for 2 hours. Dilute with ethyl acetate (60mL) and wash with 1N hydrochloric acid (30mL) and saturated sodium hydrogen carbonate (2X30mL). Dry (MgSO<sub>4</sub>), evaporate the solvent in vacuo and purify by silica gel chromatography to give [6α(R\*), 11bβ]-6-[(S)-(5-(4-methoxybenzylthio)-5-oxo-2-(carbo-t-butyloxy)-2,4,5,6-tetrahydro-cyclopenta[c]pyrrole)-methylamino]-1,2,3,5,6,7,11b-heptahydro-5-oxo-pyrrolo[2,1-a][2]benzazepine-3(S)-carboxylic acid, methyl ester.

Dissolve [6α(R\*), 11bβ]-6-[(S)-(5-(4-methoxybenzylthio)-5-oxo-2-(carbo-t-butyloxy)-2,4,5,6-tetrahydro-cyclopenta[c]pyrrole)methylamino]-1,2,3,5,6,7,11b-heptahydro-5-oxo-pyrrolo[2,1-a][2]benzazepine-3(S)-carboxylic acid, methyl ester (105mg, 0.163mmol) in methylene chloride (3mL) and cool to 0°C. Treat with trifluoroacetic acid (1.5mL), anisole (0.19mL, 1.7mmol), and mercuric acetate (65mg, 0.2mmol). Stir at 0°C for 3 hours then bubble hydrogen sulfide gas through the solution for 10 minutes. Filter and wash with methylene chloride. Wash organic phase with water (20mL), dry (Na<sub>2</sub>SO<sub>4</sub>) and evaporate the solvent in vacuo. Purify by silica gel chromatography to give the title compound.

#### Example 19

Preparation of [6α(R\*), 11bβ]-6-[(S)-(5-thio-5-oxo-2,4,5,6-tetrahydro-cyclopenta[c]pyrrole)methylamino]-1,2,3,5,6,7,11b-heptahydro-5-oxo-pyrrolo[2,1-a][2]benzazepine-3(S)-carboxylic acid

Dissolve [6α(R\*), 11bβ]-6-[(S)-(5-Thio-5-oxo-2,4,5,6-tetrahydro-cyclopenta[c]pyrrole)methylamino]-1,2,3,5,6,7,11b-heptahydro-5-oxo-pyrrolo[2,1-a][2]benzazepine-3(S)-carboxylic acid, methyl ester (42mg, 0.098mmol) in methanol (1.5mL) and degass at 0°C. Add aqueous lithium hydroxide (0.6mL of a 1N degassed solution, 0.6mmol) at 0°C. Add tetrahydrofuran to obtain solution (4mL) and stir for 17 hours at room temperature, cool in an ice bath and add 1N hydrochloric acid (1mL). Partition between methylene chloride (30mL) and water (15mL) and separate the organic phase. Dry (Na<sub>2</sub>SO<sub>4</sub>), evaporate the solvent in vacuo and purify by silica gel chromatography to give the title compound.

In a further embodiment, the present invention provides a compound of Formula I for use in inhibiting enkephalinase in a patient in need thereof comprising administering to said patient an effective enkephalinase inhibitory amount of a compound of Formula (I).

As used herein, the term "patient" refers to warm-blooded animals or mammals, including mice, rats and humans. A patient is in need of treatment to inhibit enkephalinase when the patient is suffering from acute or chronic pain and is in need of an endorphin- or enkephalin-mediated analgesic effect. In addition, a patient is in need of treatment to inhibit enkephalinase when the patient is suffering from a disease state characterized by abnormalities in fluid, electrolyte, blood pressure, intraocular pressure, renin, or aldosterone homeostasis, such as, but not limited to, hypertension, renal diseases, hyperaldosteronemia, cardiac hypertrophy, glaucoma and congestive heart failure. In these instances the patient is in need of an ANP-mediated diuretic, natriuretic, hypotensive, hypoaldosteronemic effect. Inhibition of enkephalinase would provide an endorphin- or enkephalin-mediated analgesic effect by inhibiting the metabolic degradation of endorphins and enkephalins. Inhibition of enkephalinase would provide an ANP-mediated diuretic, natriuretic, hypotensive, hypoaldosteronemic effect by inhibiting the metabolic degradation of ANP.

In addition, a patient is in need of treatment to inhibit enkephalinase when the patient is in need of an antidepressant effect or a reduction in severity of withdrawal symptoms associated with termination of opiate or morphine administration.

The identification of those patients who are in need of treatment to inhibit enkephalinase is well within the ability and knowledge of one skilled in the art. A clinician skilled in the art can readily identify, by the use of clinical tests, physical examination and medical/family history, those patients who are in need of an endorphin- or enkephalin-mediated analgesic effect or who are in need of an ANP-mediated diuretic, natriuretic, hypotensive or hypoaldosteronemic effect.

An effective enkephalinase inhibitory amount of a compound of Formula (I) is an amount which is effective in inhibiting enkephalinase and in thus inhibiting the metabolic degradation of the naturally-occurring circulating regulatory peptides such as the endorphins, including enkephalins, and ANP. Successful treatment is also understood to include prophylaxis in treating a patient in those instances such as, for example, in a pre-operative procedure, where a patient will be suffering from acute or chronic pain in the near future.

An effective enkephalinase inhibitory amount of a compound of Formula (I) is an amount which is effective in inhibiting enkephalinase in a patient in need thereof which results, for example, in endorphin- or enkephalin-mediated analgesic effects or in ANP-mediated diuretic, natriuretic, hypotensive, hypoaldosteronemic effect.

An effective enkephalinase inhibitory dose can be readily determined by the use of conventional techniques and by observing results obtained under analogous circumstances. In determining the effective dose, a number of factors are considered including, but not limited to: the species of patient; its size, age, and general health; the specific disease involved; the degree of or involvement or the severity of the disease; the response of the individual patient; the particular compound administered; the mode of administration; the bioavailability characteristics of the preparation administered; the dose regimen selected; and the use of concomitant medication.

An effective enkephalinase inhibitory amount of a compound of Formula (I) will generally vary from about 0.01 milligram per kilogram of body weight per day (mg/kg/day) to about 20 mg/kg/day. A daily dose of from about 0.1 mg/kg to about 10 mg/kg is preferred.

In addition, the present invention further provides a compound of Formula I for use in inhibiting ACE in a patient in need thereof comprising administering to said patient an effective ACE inhibitory amount of a compound of Formula (I). A patient is in need of treatment to inhibit ACE when the patient is suffering from hypertension, chronic congestive heart failure, hyperaldosteronemia or cognitive disorders. Inhibition of ACE reduces levels of angiotensin II and thus inhibits the vasopressor, hypertensive and hyperaldosteronemic effects caused thereby. An effective ACE inhibitory amount of a compound of Formula (I) is that amount which is effective in inhibiting ACE in a patient in need thereof which results, for example, in a hypotensive effect. An effective ACE inhibitory amount and an effective ACE inhibitory dose are the same as that described above for an effective enkephalinase inhibitory amount and dose.

In effecting treatment of a patient, compounds of Formula (I) can be administered in any form or mode which makes the compound bioavailable in effective amounts, including oral and parenteral routes. For example, the compound can be administered orally, subcutaneously, intramuscularly, intravenously, transdermally, intranasally, rectally, and the like. Oral administration is generally preferred. One skilled in the art of preparing Formulations can readily select the proper form and mode of administration depending upon the disease state to be treated, the stage of the disease, and other relevant circumstances.

Compounds of Formula (I) can be administered in the form of pharmaceutical compositions or medicaments which are made by combining the compounds of Formula (I) with pharmaceutically acceptable carriers or excipients, the proportion and nature of which are determined by the chosen route of administration, and standard pharmaceutical practice.

In another embodiment, the present invention provides compositions comprising a compound of Formula (I) in admixture or otherwise in association with one or more inert carriers. These compositions are useful, for example, as assay standards, as convenient means of making bulk shipments, or as pharmaceutical compositions. An assayable amount of a compound of Formula (I) is an amount which is readily measurable by standard assay procedures and techniques as are well known and appreciated by those skilled in the art. Assayable amounts of a compound of Formula (I) will generally vary from about 0.001% to about 75% of the composition by weight. Inert carriers can be any material which does not degrade or otherwise covalently react with a compound of Formula (I). Examples of suitable inert carriers are water; aqueous buffers, such as those which are generally useful in High Performance Liquid Chromatography (HPLC) analysis; organic solvents, such as acetonitrile, ethyl acetate, hexane and the like; and pharmaceutically acceptable carriers or excipients.

More particularly, the present invention provides pharmaceutical compositions comprising an effective amount of a compound of Formula (I) in admixture or otherwise in association with one or more pharmaceutically acceptable carriers or excipients.

The pharmaceutical compositions or medicaments are prepared in a manner well known in the pharmaceutical art. The carrier or excipient may be a solid, semi-solid, or liquid material which can serve as a vehicle or medium for the active ingredient. Suitable carriers or excipients are well known in the art. The pharmaceutical composition may be adapted for oral or parenteral use and may be administered to the patient in the form of tablets, capsules, suppositories, solution, suspensions, or the like.

The pharmaceutical compositions may be administered orally, for example, with an inert diluent or with an edible carrier. They may be enclosed in gelatin capsules or compressed into tablets. For the purpose of oral therapeutic administration, the compounds of Formula (I) may be incorporated with excipients and used in the form of tablets, troches, capsules, elixirs, suspensions, syrups, wafers, chewing gums and the like. These preparations should contain at least 4% of the compound of Formula (I), the active ingredient, but may be varied depending upon the particular form and may conveniently be between 4% to about 70% of the weight of the unit. The amount of the active ingredient present in compositions is such that a unit dosage form suitable for administration will be obtained.

The tablets, pills, capsules, troches and the like may also contain one or more of the following adjuvants: binders, such as microcrystalline cellulose, gum tragacanth or gelatin; excipients, such as starch

or lactose, disintegrating agents such as alginic acid, Primogel, corn starch and the like; lubricants, such as magnesium stearate or Sterotex; glidants, such as colloidal silicon dioxide; and sweetening agents, such as sucrose or saccharin may be added or flavoring agents, such as peppermint, methyl salicylate or orange flavoring. When the dosage unit form is a capsule, it may contain, in addition to materials of the above type, a liquid carrier such as polyethylene glycol or a fatty oil. Other dosage unit forms may contain other various materials which modify the physical form of the dosage unit, for example, as coatings. Thus, tablets or pills may be coated with sugar, shellac, or other enteric coating agents. A syrup may contain, in addition to the active ingredient, sucrose as a sweetening agent and certain preservatives, dyes and colorings and flavors. Materials used in preparing these various compositions should be pharmaceutically pure and non-toxic in the amounts used.

For the purpose of parenteral administration, the compounds of Formula (I) may be incorporated into a solution or suspension. These preparations should contain at least 0.1% of a compound of the invention, but may be varied to be between 0.1 and about 50% of the weight thereof. The amount of the active ingredient present in such compositions is such that a suitable dosage will be obtained.

The solutions or suspensions may also include one or more of the following adjuvants: sterile diluents such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerine, propylene glycol or other synthetic solvents; antibacterial agents such as benzyl alcohol or methyl paraben; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylene diaminetetraacetic acid; buffers such as acetates, citrates or phosphates and agents for the adjustment of toxicity such as sodium chloride or dextrose. The parenteral preparation can be enclosed in ampules, disposable syringes or multiple dose vials made of glass or plastic.

As with any group of structurally related compounds which possess a particular generic utility, certain groups and configurations are preferred for compounds of Formula (I) in their end-use application.

The compounds of Formula (1) wherein  $B_1$  is hydrogen or alkoxy are preferred. Compounds of Formula (1) wherein  $R_1$  and  $R_3$  are hydrogen and  $n$  is 0 are preferred.

It is, of course, understood that the compounds of Formula (I) may exist in a variety of isomeric configurations including structural as well as stereo isomers. It is further understood that the present invention encompasses those compounds of Formula (I) in each of their various structural and stereo isomeric configurations as individual isomers and as mixtures of isomers.

The following specific compounds of Formula (1) are particularly preferred in the end-use application of the compounds of the present invention:

[4S-[4 $\alpha$ , 7 $\alpha$ (R\*), 12b $\beta$ ]]-7-[(2-Thio-2-oxoindan)methylamino]-1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido[2,1-a]-[2]benzazepine-4-carboxylic acid;  
[4S-[4 $\alpha$ , 7 $\alpha$ (R\*), 12b $\beta$ ]]-7-[(2-mercaptomethyl-2-oxoindan)methylamino]-1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido[2,1-a]-[2]benzazepine-4-carboxylic acid;  
[4S-[4 $\alpha$ , 7 $\alpha$ (R\*), 12b $\beta$ ]]-7-[(2-(2-mercaptoethyl)-2-oxoindan)methylamino]-1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido[2,1-a]-[2]benzazepine-4-carboxylic acid;  
[4S-[4 $\alpha$ , 7 $\alpha$ (R\*), 12b $\beta$ ]]-7-[(1-Thio-1-oxocyclopentane)methylamino]-1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido[2,1-a]-[2]benzazepine-4-carboxylic acid.

The following studies illustrate the utility of the compounds of the present invention as enkephalinase inhibitors and as ACE inhibitors.

Enkephalinase is partially purified from rat kidney. The enzyme is extracted from the microvilli fraction by using Triton X-100 according to the method of Malfroy and Schwartz [*J.Biol.Chem.* 259, 14365-14370 (1984)] or by using a proteolytic treatment according to the method of Almenoff and Orlowski [*Biochem.* 22, 590-599 (1983)]. The enzyme is further purified by anion exchange chromatography (Mono Q™ column, Pharmacia) using a Pharmacia FPLC system. The enzyme activity may be measured by the fluorometric methods of Florentin et al. [*Anal.Biochem.* 141, 62-69 (1984)] or of Almenoff and Orlowski [*J.Neurochemistry* 42, 151-157 (1984)]. The enzyme is assayed in 50mM HEPES buffer (pH 7.4) in a 3.0 mL reaction volume containing 12  $\mu$ M of the substrate dansyl-D-AlaGly(p-nitro)PheGly ( $K_m$  = 40 $\mu$ M) at 25°C. The substrate (and inhibitor) is added from a concentrated stock solution in DMSO (up to 0.1 mL DMSO final volume). The enzyme in a small volume (approximately 0.1  $\mu$ g of FPLC purified protein) is added to initiate the reaction and the rate of fluorescence increase is recorded continuously using a fluorometer (excitation at 339nm, emission at 562nm).

The enzymatic activity of ACE is monitored using the spectrophotometric substrate described by Holmquist et al. [*Anal.Biochem.* 95, 540-548 (1979)] and the buffer system described by Ryan [*Methods of Enzymatic Analysis*, 3rd ed., H. U. Bergmeyer, editor; vol. V, Verlag Chemie, Weinheim, 1983, pp. 20-34].

The results of the analysis of enzymatic activity as described in Table 1 indicate that the compounds of the present invention are inhibitors of enkephalinase as well as inhibitors of ACE.


Table 1

K <sub>i</sub> 's of Compounds of Formula (I) as Inhibitors of Enkephalinase and of ACE		
Compound of Formula (I)	ACE, K <sub>i</sub> (nM)	ENK, K <sub>i</sub> (nM)
102,708	<45	<0.2
100,346	<500	<0.2
100,273	>250	0.8
102,769	<7	<0.2
101,069	>7	4

102,708 = [4S-[4 $\alpha$ , 7 $\alpha$ (R<sup>\*</sup>), 12b $\beta$ ]]-7-[(2-Thio-2-oxoindan)methylamino]-1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido[2,1-a][2]benzazepine-4-carboxylic acid  
 100,346 = [4S-[4 $\alpha$ , 7 $\alpha$ (R<sup>\*</sup>), 12b $\beta$ ]]-7-[(2-mercaptomethyl-2-oxoindan)methylamino]-1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido[2,1-a][2]benzazepine-4-carboxylic acid  
 100,273 = [4S-[4 $\alpha$ , 7 $\alpha$ (R<sup>\*</sup>), 12b $\beta$ ]]-7-[(2-(2-mercaptoethyl)-2-oxoindan)methylamino]-1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido[2,1-a][2]benzazepine-4-carboxylic acid  
 102,769 = [4S-[4 $\alpha$ , 7 $\alpha$ (R<sup>\*</sup>), 12b $\beta$ ]]-7-[(1-Thio-1-oxocyclopentane)methylamino]-1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido[2,1-a][2]benzazepine-4-carboxylic acid  
 101,069 = [4S-[4 $\alpha$ , 7 $\alpha$ (R<sup>\*</sup>), 12b $\beta$ ]]-7-[(2-Thio-1-oxo-2-methyl)-2-propylamino]-1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido[2,1-a][2]benzazepine-4-carboxylic acid

## Claims

1. A compound of the formula



wherein

B<sub>1</sub> and B<sub>2</sub> are each independently hydrogen; hydroxy; -OR<sub>2</sub> wherein R<sub>2</sub> is a C<sub>1</sub>-C<sub>4</sub> alkyl or an Ar-Y group wherein Ar is aryl and Y is a hydrogen or C<sub>1</sub>-C<sub>4</sub> alkyl; or, where B<sub>1</sub> and B<sub>2</sub> are attached to adjacent carbon atoms, B<sub>1</sub> and B<sub>2</sub> can be taken together with said adjacent carbons to form a benzene ring or methylenedioxy;


A is a bond, methylene, oxygen, sulfur, NR<sub>4</sub> or NCOR<sub>5</sub> wherein R<sub>4</sub> is hydrogen, a C<sub>1</sub>-C<sub>4</sub> alkyl or an Ar-Y- group and R<sub>5</sub> is -CF<sub>3</sub>, a C<sub>1</sub>-C<sub>10</sub> alkyl or an Ar-Y- group;

R<sub>1</sub> is hydrogen, acetyl, -CH<sub>2</sub>OC(O)C(CH<sub>3</sub>)<sub>3</sub> or benzoyl;

R<sub>3</sub> is hydrogen or -CH<sub>2</sub>OC(O)C(CH<sub>3</sub>)<sub>3</sub>;


n is an integer 0 to 3; and

Q is a group of the formula



wherein Z is O, NH or S; and m is an integer 1 to 5.

2. A compound according to Claim 1 wherein Q is



3. A compound according to Claim 2 wherein A is methylene.

4. A compound according to Claim 3 wherein R<sub>1</sub> and R<sub>3</sub> are hydrogen.

5. A compound according to Claim 4 wherein n is 0.

6. A compound according to Claim 4 wherein n is 1.

7. A compound according to Claim 4 wherein n is 2.

8. A compound according to Claim 1 wherein Q is



9. A compound according to Claim 8 wherein A is methylene.

10. A compound according to Claim 9 wherein R<sub>1</sub> and R<sub>3</sub> are hydrogen.

11. A compound according to Claim 10 wherein n is 0.

12. A compound according to Claim 1 wherein A is -O-.

13. A compound according to Claim 1 wherein A is -S-.

14. A compound according to Claim 1 wherein A is -NH-.

15. A compound of Claim 1 wherein the compound is [4S-[4 $\alpha$ , 7 $\alpha$ (R'), 12b $\beta$ ]]-7-[(2-Thio-2-oxoindan)-methylamino]-1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido[2,1-a][2]benzazepine-4-carboxylic acid;

[4S-[4 $\alpha$ , 7 $\alpha$ (R'), 12b $\beta$ ]]-7-[(2-mercaptomethyl-2-oxoindan)methylamino]-1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido[2,1-a][2]benzazepine-4-carboxylic acid;

[4S-[4 $\alpha$ , 7 $\alpha$ (R'), 12b $\beta$ ]]-7-[(2-(2-mercaptoethyl)-2-oxoindan)methylamino]-1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido[2,1-a][2]benzazepine-4-carboxylic acid;

[4S-[4 $\alpha$ , 7 $\alpha$ (R'), 12b $\beta$ ]]-7-[(1-Thio-1-oxocyclopentane)methylamino]-1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido[2,1-a][2]benzazepine-4-carboxylic acid;

[4S-[4 $\alpha$ , 7 $\alpha$ (R'), 12b $\beta$ ]]-7-[(1-Thio-1-oxocyclohexane)methylamino]-1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido[2,1-a][2]benzazepine-4-carboxylic acid; or

[4S-[4 $\alpha$ , 7 $\alpha$ (R'), 12b $\beta$ ]]-7-[(1-Thio-1-oxocyclopropane)methylamino]-1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido[2,1-a][2]benzazepine-4-carboxylic acid; or

[4S-[4 $\alpha$ , 7 $\alpha$ (R'), 12b $\beta$ ]]-7-[(2-Thio-1-oxo-2-methyl)-2-propylamino]-1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido[2,1-a][2]benzazepine-4-carboxylic acid.

16. A compound according to any of Claims 1 to 15 for use as a pharmaceutically active agent.

17. A compound according to Claim 16 for use in inhibiting enkephalinase.

18. A compound according to Claim 17 for use as an endorphin- or enkephalin-mediated analgesic agent.

19. A compound according to Claim 17 for use as an ANP-mediated hypotensive agent.

20. A compound according to Claim 17 for use as an ANP-mediated diuretic agent.

21. A compound according to Claim 17 for use in treating congestive heart failure.

22. A compound according to Claim 16 for use in inhibiting ACE.

23. A compound according to Claim 22 for use as a hypotensive agent.

24. A compound according to Claim 22 for use as a cognition enhancing agent.


25. A compound according to Claim 22 for use in treating congestive heart failure.

26. A composition comprising an assayable amount of a compound according to any of Claims 1 to 15 in admixture or otherwise in association with an inert carrier.

27. A pharmaceutical composition comprising an effective immunosuppressive amount of a compound according to any of Claims 1 to 15 in admixture or otherwise in association with one or more pharmaceutically acceptable carriers or excipients.

**Claims for the following Contracting State : ES**

1. A process for preparing the compounds of formula



wherein

B<sub>1</sub> and B<sub>2</sub> are each independently hydrogen; hydroxy; -OR<sub>2</sub> wherein R<sub>2</sub> is a C<sub>1</sub>-C<sub>4</sub> alkyl or an Ar-Y group wherein Ar is aryl and Y is a hydrogen or C<sub>1</sub>-C<sub>4</sub> alkyl; or, where B<sub>1</sub> and B<sub>2</sub> are attached to adjacent carbon atoms, B<sub>1</sub> and B<sub>2</sub> can be taken together with said adjacent carbons to form a benzene ring or methylenedioxy;

A is a bond, methylene, oxygen, sulfur, NR<sub>4</sub> or NCOR<sub>5</sub> wherein R<sub>4</sub> is hydrogen, a C<sub>1</sub>-C<sub>4</sub> alkyl or an Ar-Y group and


R<sub>5</sub> is -CF<sub>3</sub>, a C<sub>1</sub>-C<sub>10</sub> alkyl or an Ar-Y group;

R<sub>1</sub> is hydrogen;

R<sub>3</sub> is hydrogen;


n is an integer 0 to 3; and

Q is a group of the formula




wherein Z is O, NH or S; and m is an integer 1 to 5, comprising reacting a compound of the formula





wherein  $B_1$ ,  $B_2$ , A, n and Q are as defined above and Pg is a suitable protecting group with mercuric acetate in the presence of an acid.

2. A process for preparing the compounds of formula



wherein

$B_1$  and  $B_2$  are each independently hydrogen; hydroxy;  $-OR_2$  wherein  $R_2$  is a  $C_1$ - $C_4$  alkyl or an Ar-Y group wherein Ar is aryl and Y is a hydrogen or  $C_1$ - $C_4$  alkyl; or, where  $B_1$  and  $B_2$  are attached to adjacent carbon atoms,  $B_1$  and  $B_2$  can be taken together with said adjacent carbons to form a benzene ring or methylenedioxy;

A is a bond, methylene, oxygen, sulfur,  $NR_4$  or  $NCOR_5$  wherein  $R_4$  is hydrogen, a  $C_1$ - $C_4$  alkyl or an Ar-Y group and

$R_5$  is  $-CF_3$ , a  $C_1$ - $C_{10}$  alkyl or an Ar-Y group;

$R_1$  is  $-CH_2OC(O)C(CH_3)_3$ ;

$R_3$  is hydrogen;

n is an integer 0 to 3; and

Q is a group of the formula



20



35

## 40



59

•

A is a bond, methylene, oxygen, sulfur,  $\text{NR}_4$  or  $\text{NCOR}_5$  wherein  $\text{R}_4$  is hydrogen, a  $\text{C}_1\text{-C}_4$  alkyl or an Ar-Y group and


$\text{R}_5$  is  $-\text{CF}_3$ , a  $\text{C}_1\text{-C}_{10}$  alkyl or an Ar-Y group;

$\text{R}_1$  is acetyl or benzoyl;


$\text{R}_3$  is hydrogen;

n is an integer 0 to 3; and

Q is a group of the formula




wherein Z is O, NH or S; and m is an integer 1 to 5, comprising reacting a compound of the formula



wherein  $\text{B}_1$ ,  $\text{B}_2$ , A, n and Q are as defined above,  $\text{R}_1$  and  $\text{R}_3$  are hydrogen with an acylating agent.

4. A process for preparing the compounds of formula



wherein

B<sub>1</sub> and B<sub>2</sub> are each independently hydrogen; hydroxy; -OR<sub>2</sub> wherein R<sub>2</sub> is a C<sub>1</sub>-C<sub>4</sub> alkyl or an Ar-Y group wherein Ar is aryl and Y is a hydrogen or C<sub>1</sub>-C<sub>4</sub> alkyl; or, where B<sub>1</sub> and B<sub>2</sub> are attached to adjacent carbon atoms, B<sub>1</sub> and B<sub>2</sub> can be taken together with said adjacent carbons to form a benzene ring or methylenedioxy;

A is a bond, methylene, oxygen, sulfur, NR<sub>4</sub> or NCOR<sub>5</sub> wherein R<sub>4</sub> is hydrogen, a C<sub>1</sub>-C<sub>4</sub> alkyl or an Ar-Y group and


R<sub>5</sub> is -CF<sub>3</sub>, a C<sub>1</sub>-C<sub>10</sub> alkyl or an Ar-Y group;

R<sub>1</sub> is acetyl or benzoyl;


R<sub>3</sub> is -CH<sub>2</sub>OC(O)C(CH<sub>3</sub>)<sub>3</sub>;

n is an integer 0 to 3; and

Q is a group of the formula




wherein Z is O, NH or S; and m is an integer 1 to 5, comprising reacting a compound of the formula



wherein  $B_1$ ,  $B_2$ , A,  $R_1$ , n and Q are as defined above and  $R_3$  is hydrogen with chloromethyl pivalate in the presence of a suitable base.

5. A process for preparing the compounds of formula



wherein

$B_1$  and  $B_2$  are each independently hydrogen; hydroxy;  $-OR_2$  wherein  $R_2$  is a  $C_1$ - $C_4$  alkyl or an Ar-Y group wherein Ar is aryl and Y is a hydrogen or  $C_1$ - $C_4$  alkyl; or, where  $B_1$  and  $B_2$  are attached to adjacent carbon atoms,  $B_1$  and  $B_2$  can be taken together with said adjacent carbons to form a benzene ring or methylenedioxy;

A is a bond, methylene, oxygen, sulfur,  $NR_4$  or  $NCOR_5$  wherein  $R_4$  is hydrogen, a  $C_1$ - $C_4$  alkyl or an Ar-Y group and


$R_5$  is  $-CF_3$ , a  $C_1$ - $C_{10}$  alkyl or an Ar-Y group;

$R_1$  is hydrogen;


$R_3$  is  $-CH_2OC(O)C(CH_3)_3$ ;

n is an integer 0 to 3; and

Q is a group of the formula




15 wherein Z is O, NH or S; and m is an integer 1 to 5, comprising reacting a compound of the formula



wherein B<sub>1</sub>, B<sub>2</sub>, A, R<sub>3</sub>, n and Q are as defined above and R<sub>1</sub> is acetyl or benzoyl with ammonia.

35 6. A process for preparing the compounds of formula



wherein

55 B<sub>1</sub> and B<sub>2</sub> are each independently hydrogen; hydroxy; -OR<sub>2</sub> wherein R<sub>2</sub> is a C<sub>1</sub>-C<sub>4</sub> alkyl or an Ar-Y group wherein Ar is aryl and Y is a hydrogen or C<sub>1</sub>-C<sub>4</sub> alkyl; or, where B<sub>1</sub> and B<sub>2</sub> are attached to adjacent carbon atoms, B<sub>1</sub> and B<sub>2</sub> can be taken together with said adjacent carbons to form a benzene ring or methylenedioxy;

A is a bond, methylene, oxygen, sulfur, NR<sub>4</sub> or NCOR<sub>5</sub> wherein R<sub>4</sub> is hydrogen, a C<sub>1</sub>-C<sub>4</sub> alkyl or an

Ar-Y group and


R<sub>5</sub> is -CF<sub>3</sub>, a C<sub>1</sub>-C<sub>10</sub> alkyl or an Ar-Y group;

R<sub>1</sub> is -CH<sub>2</sub>OC(O)C(CH<sub>3</sub>)<sub>3</sub>;


R<sub>3</sub> is -CH<sub>2</sub>OC(O)C(CH<sub>3</sub>)<sub>3</sub>;

n is an integer 0 to 3; and

Q is a group of the formula




wherein Z is O, NH or S; and m is an integer 1 to 5, comprising reacting a compound of the formula




wherein B<sub>1</sub>, B<sub>2</sub>, A, n and Q are as defined above and R<sub>1</sub> and R<sub>3</sub> are hydrogen with chloromethyl pivalate in the presence of a suitable base.

7. A process according to any of Claims 1 to 6 wherein Q is



8. A process according to Claim 7 wherein A is methylene.

9. A process according to Claim 1 wherein R<sub>1</sub> and R<sub>3</sub> are hydrogen, Q is



and A is methylene.

10. A process according to Claim 9 wherein n is 0.

11. A process according to Claim 9 wherein n is 1.

12. A process according to Claim 9 wherein n is 2.

13. A process according to any of Claims 1 to 6 wherein Q is



14. A process according to Claim 13 wherein A is methylene.

15. A process according to Claim 1 wherein R<sub>1</sub> and R<sub>3</sub> are hydrogen, Q is



and A is methylene.

16. A process according to Claim 15 wherein n is 0.

17. A process according to any of Claims 1 to 6 wherein A is -O-.

18. A process according to any of Claims 1 to 6 wherein A is -S-.

19. A process according to any of Claims 1 to 6 wherein A is -NH-.

20. A process according to any of Claims 1 to 6 wherein the compound to be produced is [4S-[4 $\alpha$ , 7 $\alpha$ (R\*), 12b $\beta$ ]]-7-[(2-Thio-2-oxoindan)methylamino]-1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido[2,1-a][2]-benzazepine-4-carboxylic acid;

4S-[4 $\alpha$ , 7 $\alpha$ (R\*), 12b $\beta$ ]]-7-[(2-mercaptomethyl-2-oxoindan)methylamino]-1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido[2,1-a][2]benzazepine-4-carboxylic acid;

[4S-[4 $\alpha$ , 7 $\alpha$ (R\*), 12b $\beta$ ]]-7-[(2-(2-mercaptoethyl)-2-oxoindan)methylamino]-1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido[2,1-a][2]benzazepine-4-carboxylic acid;

[4S-[4 $\alpha$ , 7 $\alpha$ (R\*), 12b $\beta$ ]]-7-[(1-Thio-1-oxocyclopentane)methylamino]-1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido[2,1-a][2]benzazepine-4-carboxylic acid;

[4S-[4 $\alpha$ , 7 $\alpha$ (R\*), 12b $\beta$ ]]-7-[(1-Thio-1-oxocyclohexane)methylamino]-1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido[2,1-a][2]benzazepine-4-carboxylic acid;



[4S-[4 $\alpha$ , 7 $\alpha$ (R), 12b $\beta$ ]]-7-[(1-Thio-1-oxocyclopropane)methylamino]-1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido[2,1-a][2]benzazepine-4-carboxylic acid; or

[4S-[4 $\alpha$ , 7 $\alpha$ (R), 12b $\beta$ ]]-7-[(2-Thio-1-oxo-2-methyl)-2-propylamino]-1,2,3,4,6,7,8,12b-octahydro-6-oxopyrido[2,1-a][2]benzazepine-4-carboxylic acid.

21. A composition for nonpharmaceutical purposes comprising an assayable amount of a compound obtainable according to the process of any of Claims 1 to 20 in admixture or otherwise in association with an inert carrier.

22. A process for producing a pharmaceutical composition comprising mixing an effective immunosuppressive amount of a compound obtainable according to the process of any of Claims 1 to 20 with one or more pharmaceutically acceptable carriers or excipients.